

Nancy Kinoti

DMA TRAINING
LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
 See slide 65 in DMA Packette

Mental Illness Anxiety Disorders	Inhalers Allergy / Asthma	Hyperlipidemia Statins	Diabetes Endocrine & Metabolic
Abilify Ativan Clozaril Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa	Advair Discus Atrovent Flonase Flovent Loratadine Proventil	Crestor Lipitor Zocor	Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid
Seizures	Gastrointestinal Disorder Constipation	Blood Pressure Meds	Pain & Inflammation
Dilantin Keppra Neurontin Topamax	Colace Miralax Prilosec Protonix Zantac	HCTZ (hydrochlorothiazide) Lisinopril Toprol Tenormin	Flexeril Motrin Norco Tylenol with Codeine Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

Nancy Ninoti

DRUG NAME	Lamictal
GENERIC NAME	Lamotrigine
DOSAGE RANGE	25mg, 100mg, 150mg, 200mg; 25mg, 50mg, 100mg, 200mg
HOW TO TAKE	Swallow the tablet whole and do not crush, chew or break
USES	bipolar I disorder, Partial seizures, seizures, headache
SIDE EFFECTS	vertigo, vomiting, blurred vision
WARNINGS	serious rash

DRUG NAME	Zyprexa
GENERIC NAME	Olanzapine
DOSAGE RANGE	2.5mg, 5mg, 7.5mg, 10mg, 15mg, 20mg, 10, 5mg
HOW TO TAKE	Olanzapine is injected into a muscle
USES	Schizophrenia, bipolar I disorder, agitation
SIDE EFFECTS	weight gain, insomnia, back pain, tremor
WARNINGS	Dementia related psychosis

DRUG NAME	Risperdal
GENERIC NAME	Risperidone
DOSAGE RANGE	0.25, 0.5, 1mg, 1mg, 2mg, 3mg, 4mg, Sol: 1mg per ml
HOW TO TAKE	Place the tablet in your mouth and allow it to dissolve
USES	Schizophrenia, bipolar I disorder, Tourette Syndrome
SIDE EFFECTS	Fever, cough, anxiety, tremor, headache
WARNINGS	Dementia-Related psychosis not approved

DRUG NAME	Klonopin
GENERIC NAME	Clonazepam
DOSAGE RANGE	Tab 0.5mg, 1mg, 2mg
HOW TO TAKE	use dry hands to remove the tablet & place in your mouth
USES	Seizure disorder, Panic disorder, anxiety, restless leg
SIDE EFFECTS	fatigue, confusion, irritability, rash, constipation
WARNINGS	concomitant benzodiazepine use w/ opioids may result in profound sedation

DRUG NAME	Seroquel
GENERIC NAME	Quetiapine
DOSAGE RANGE	25mg, 50mg, 100mg, 200mg, 300mg, 400mg
HOW TO TAKE	Swallow the tablet whole and do not swallow
USES	Schizophrenia, bipolar, generalized anxiety disorder
SIDE EFFECTS	Anemia, weight gain, tremor, constipation
WARNINGS	Dementia related psychosis, suicidality

Nancy
Kinoti

DRUG NAME	Abilify
GENERIC NAME	Aripiprazole
DOSAGE RANGE	2mg - 30mg
HOW TO TAKE	Take it by mouth, liquid or pill with/without food
USES	For schizophrenia, bipolar, major depressive
SIDE EFFECTS	Weight gain, increased appetite, headache
WARNINGS	dementia-related psychosis, suicidality

DRUG NAME	Ativan
GENERIC NAME	Lorazepam
DOSAGE RANGE	0.5mg, 1mg, 2mg; inj: 2mg per mL, 4mg per mL
HOW TO TAKE	Potently measure liquid medication carefully
USES	Anxiety, insomnia, status epilepticus, nausea, vomiting
SIDE EFFECTS	Sedation, dizziness, menstrual irregularities
WARNINGS	may result in depression, coma, or death

DRUG NAME	Depakote
GENERIC NAME	Divalproex sodium
DOSAGE RANGE	125mg, 250mg, 500mg
HOW TO TAKE	by mouth drink plenty of water while you are taking ^{the} med.
USES	Partial seizures, absence seizures, bipolar disorder, migraine
SIDE EFFECTS	headache, vomiting, diarrhea, tremor, dizziness
WARNINGS	Serious or fatal hepatic failure, risk of acute liver failure

DRUG NAME	Invega
GENERIC NAME	Paliperidone
DOSAGE RANGE	1.5mg, 3mg, 6mg, 9mg
HOW TO TAKE	Do not cut/crush/chew tab/DIC if unexplained clear
USES	Schizophrenia, Schizoaffective disorder
SIDE EFFECTS	nausea, headache, anxiety, dizziness
WARNINGS	not approved for dementia-related psychosis

DRUG NAME	Lithium
GENERIC NAME	generic
DOSAGE RANGE	cap: 150mg, 300mg, 600mg, Tab: 300mg, Er Tab 300mg, L 50mg
HOW TO TAKE	by mouth, swallow the tablet while do not crush, chew, break
USES	bipolar & disorder, maintenance tx, acute manic/mixed, schiz
SIDE EFFECTS	flone, edema, fatigue, rash, anorexia
WARNINGS	Lithium toxicity closely related to serum lithium levels and can occur at close to therapeutic levels