

DMA TRAINING

LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
See slide 65 in DMA Packette

| Mental Illness Anxiety Disorders | Inhalers Allergy / Asthma | Hyperlipidemia Statins | Diabetes Endocrine & Metabolic |
|--|--|--|---|
| Abilify Ativan Clozaril Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa | Advair Discus Atrovent Flonase Flovent Loratadine Proventil | Crestor Lipitor Zocor | Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid |
| | | | |
| Seizures | Gastrointestinal Disorder Constipation | Blood Pressure Meds | Pain & Inflammation |
| Dilantin Keppra Neurontin Topamax | Colace Miralax Prilosec Protonix Zantac | HCTZ (hydrochlorothiazide) Lisinopril Toprol Tenormin | Flexeril Motrin Norco Tylenol with Codeine Ultram |

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1

1217

DMA Code #2

1109

DMA Code #3

0917

| | |
|--------------|--|
| DRUG NAME | Ativan |
| GENERIC NAME | Lorazepam |
| DOSAGE RANGE | tab. .5mg, 1mg, 2mg. Inj. 2mg per mL 4mg per mL |
| HOW TO TAKE | By mouth or injection |
| USES | anxiety, insomnia, preoperative sedation |
| SIDE EFFECTS | Drowsiness, dizziness, blurred vision |
| WARNINGS | Risks from concomitant opioid use, addiction, abuse, misuse, dependence reaction |

| | |
|--------------|--|
| DRUG NAME | Depakote |
| GENERIC NAME | divalproex Sodium |
| DOSAGE RANGE | Or Tab. 125mg, 250mg, 500mg |
| HOW TO TAKE | By mouth |
| USES | Bi Polar disorder, Seizures, migraines |
| SIDE EFFECTS | Hepatotoxicity, pancreatitis, SIADH |
| WARNINGS | Hepatotoxicity, Fetal risk, pancreatitis |

| | |
|--------------|---|
| DRUG NAME | Haldol |
| GENERIC NAME | haloperidol |
| DOSAGE RANGE | tab. .5mg, 1mg, 2mg, 5mg, 10mg, 20mg |
| HOW TO TAKE | By mouth or injections |
| USES | Psychosis, Tourette Syndrome, agitation |
| SIDE EFFECTS | tardive dyskinesia, hypotension |
| WARNINGS | Dementia-Related psychosis |

| | |
|--------------|---|
| DRUG NAME | Invega |
| GENERIC NAME | paliperidone |
| DOSAGE RANGE | ER Tab. 1.5mg, 3mg, 6mg, 9mg |
| HOW TO TAKE | By mouth |
| USES | Schizophrenia, Schizoaffective disorder |
| SIDE EFFECTS | arrhythmia, QT prolongation |
| WARNINGS | Dementia Related psychosis |

| | |
|--------------|---|
| DRUG NAME | Klonopin |
| GENERIC NAME | clonazepam |
| DOSAGE RANGE | Tab. .5mg, 1mg, 2mg |
| HOW TO TAKE | By mouth |
| USES | Seizure disorder, panic disorder, anxiety |
| SIDE EFFECTS | respiratory depression, abuse, tachycardia |
| WARNINGS | Risks from concomitant opioid use, addiction, abuse, misuse, dependence and |

| | |
|--------------|--|
| DRUG NAME | Lithium |
| GENERIC NAME | Lithium |
| DOSAGE RANGE | Cap: 150mg, 300mg, 600mg. TAB: 300mg ER TAB: 300mg / 450mg |
| HOW TO TAKE | By mouth |
| USES | Bi Polar disorder |
| SIDE EFFECTS | Coma, Seizures, ventricular arrhythmia |
| WARNINGS | Lithium toxicity |

| | |
|--------------|---|
| DRUG NAME | Seroquel |
| GENERIC NAME | quetiapine |
| DOSAGE RANGE | TAB: 25mg, 50mg, 100mg, 200mg, 300mg, 400mg |
| HOW TO TAKE | By mouth |
| USES | Schizophrenia, Bi Polar Disorder |
| SIDE EFFECTS | Tardive dyskinesia, depression exacerbation |
| WARNINGS | Dementia-Related psychosis, suicidality |

| | |
|--------------|--|
| DRUG NAME | Risperdal |
| GENERIC NAME | Risperidone |
| DOSAGE RANGE | TAB: .25mg, .5mg, 1mg, 2mg, 3mg, 4mg SOL: 1mg per mL |
| HOW TO TAKE | By mouth |
| USES | Schizophrenia, Bi Polar disorder, Tourette Syndrome |
| SIDE EFFECTS | hypotension, syncope, tardive dyskinesia |
| WARNINGS | Dementia Related psychosis |

| | |
|--------------|---|
| DRUG NAME | Zyprexa |
| GENERIC NAME | olanzapine |
| DOSAGE RANGE | TAB: 2.5mg, 5mg, 7.5mg, 10mg, 15mg, 20mg, Inj: 5mg per mL |
| HOW TO TAKE | By mouth or injection |
| USES | Schizophrenia, Bi Polar disorder |
| SIDE EFFECTS | hypoglycemia, stroke, dystonia |
| WARNINGS | Dementia Related psychosis |

| | |
|--------------|--|
| DRUG NAME | Topamax |
| GENERIC NAME | topiramate |
| DOSAGE RANGE | Cap: 15mg, 25mg, TAB: 25mg, 50mg, 100mg, 200mg |
| HOW TO TAKE | By mouth |
| USES | Seizures |
| SIDE EFFECTS | metabolic acidosis, osteoporosis |
| WARNINGS | N/A |