

DMA TRAINING

LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
See slide 65 in DMA Packette

Mental Illness Anxiety Disorders	Inhalers Allergy / Asthma	Hyperlipidemia Statins	Diabetes Endocrine & Metabolic
Abilify Ativan Clozaril Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa	Advair Discus Atrovent Flonase Flovent Loratadine Proventil	Crestor Lipitor Zocor	Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid
Seizures	Gastrointestinal Disorder Constipation	Blood Pressure Meds	Pain & Inflammation
Dilantin Keppra Neurontin Topamax	Colace Miralax Prilosec Protonix Zantac	HCTZ (hydrochlorothiazide) Lisinopril Toprol Tenormin	Flexeril Motrin Norco Tylenol with Codeine Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 6798

DMA Code #2 1109

DMA Code #3 1217

Jayla F

DRUG NAME	Seroquel
GENERIC NAME	Quetiapine
DOSAGE RANGE	25mg, 50mg, 100mg, 200mg 200mg 300mg 400mg
HOW TO TAKE	By mouth
USES	Schizophrenia, Bipolar Disorder, (manic) Renal Dosis
SIDE EFFECTS	Rash, Dizziness, Constipation, Hypotension, Back pain
WARNINGS	Dementia-Related Psychosis / Suicidality

DRUG NAME	Ativan Ativan
GENERIC NAME	Lorazepam
DOSAGE RANGE	0.5mg, 1mg, 2mg, INJ: 2mg per mL, 4mg per mL
HOW TO TAKE	By mouth
USES	Anxiety, insomnia, status Epilepticus, Nausea / Vomiting
SIDE EFFECTS	Dizziness, confusion, irritability, fatigue, Diplopia
WARNINGS	Risk from concomitant opioid, Addiction, Abuse, misuse, dependence and withdrawal reactions.

DRUG NAME	Klonopin
GENERIC NAME	Clonazepam
DOSAGE RANGE	0.5, 1mg, 2mg
HOW TO TAKE	By mouth
USES	Seizure, panic disorder, Anxiety, Restless leg syndrome
SIDE EFFECTS	Drowsiness,
WARNINGS	Risk from concomitant opioid use, addiction, Abuse, misuse, dependence and withdrawal reactions.

DRUG NAME	Tenormin
GENERIC NAME	Atenolol
DOSAGE RANGE	25mg, 50mg, 100mg
HOW TO TAKE	By mouth
USES	Angina, Cardiovascular prevention, migraine headache
SIDE EFFECTS	hypotension, fatigue, nausea, leg pain, depression
WARNINGS	Avoid Abrupt Cessation

DRUG NAME	Norco
GENERIC NAME	Hydrocodone / Acetaminophen
DOSAGE RANGE	5mg / 325mg, 7.5mg / 325mg, 10mg / 325mg
HOW TO TAKE	By mouth
USES	Special Note, Pain, moderate - mod. severe
SIDE EFFECTS	Sedation, nausea, vomiting, Dizziness, Rash
WARNINGS	Medication Error risk, Addiction, Abuse, misuse misuse, Respiratory Depression, Accidental Ingestion

Jayla Finch

DRUG NAME	Miralax
GENERIC NAME	Polyethylene glycol 3350
DOSAGE RANGE	<17yo. and older 20kg 17under less than 20kg
HOW TO TAKE	By mouth
USES	Osmotic laxative for relieving constipation.
SIDE EFFECTS	Nausea, Abdominal distension, cramping, diarrhea.
WARNINGS	hypersens. to drug/class/compon, caution in elderly pts, caution in renal impairment.

DRUG NAME	Lantus
GENERIC NAME	insulin Glargine
DOSAGE RANGE	INJ (Solostar U-100 pen) 100 units per mL ^{FNJ (U-100) 100 units per mL}
HOW TO TAKE	Injection
USES	Diabetes Mellitus Type 1, Type 2
SIDE EFFECTS	hypoglycemia, inj site Rxn, myalgia, Rash, headache
WARNINGS	hypersens to drug/class/compon, hypoglycemia, caution if infection, illness, or stress.

DRUG NAME	Atrivent
GENERIC NAME	ipratropium Bromide inhaled
DOSAGE RANGE	MDI: 17mcg per actuation
HOW TO TAKE	By mouth
USES	COPD, maintenance tx, Asthma exacerbation, mod-severe
SIDE EFFECTS	Bronchitis, dyspnea, nausea, sinusitis, UTI, Back pain
WARNINGS	Hypersens to drug/class/compon, Hypersens to Atropine
CAUTION IF	glaucoma, angle-closure, prostatic hypertrophy.

DRUG NAME	Lipitor
GENERIC NAME	Atorvastatin
DOSAGE RANGE	10mg, 20mg, 40mg, 80mg.
HOW TO TAKE	By mouth
USES	Hypercholesterolemia, mixed dyslipidemia
SIDE EFFECTS	URI, headache, UTI, nausea, muscle spasms
WARNINGS	Pregnancy (non-high risk pts), myopathy, Hepatic.

DRUG NAME	reaction Pantoprazole Protonix
GENERIC NAME	Cyclobenzaprine pantoprazole
DOSAGE RANGE	OR TAB: 20mg, 40mg, DR Granule 40mg per pkt INJ vial
HOW TO TAKE	By mouth By Mouth, INJ
USES	GERD, hypersecretory conditions, upper GI Bleeding
SIDE EFFECTS	Headache, nausea, vomiting, dizziness, diarrhea
WARNINGS	Hypersens to drug/class/compon, caution if long term use, high dose use, hypomagnesemia.