

Helen
Bates

DMA TRAINING
LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
See slide 65 in DMA Packette

| Mental Illness Anxiety Disorders | Inhalers Allergy / Asthma | Hyperlipidemia Statins | Diabetes Endocrine & Metabolic |
|--|--|--|---|
| Abilify Ativan Clozaryl Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa | Advair Discus Atrovent Flonase Flovent Loratadine Proventil | Crestor Lipitor Zocor | Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid |
| | | | |
| Seizures | Gastrointestinal Disorder Constipation | Blood Pressure Meds | Pain & Inflammation |
| Dilantin Keppra Neurontin Topamax | Colace Miralax Prilosec Protonix Zantac | HCTZ (hydrochlorothiazide) Lisinopril Toprol Tenormin | Flexeril Motrin Norco Tylenol with Codeine Ultram |

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 7821

DMA Code #2 2221

DMA Code #3 3482

| | |
|--------------|--|
| DRUG NAME | Ability |
| GENERIC NAME | Dripiprazole |
| DOSAGE RANGE | 2.5-10-15 mg PO qd - Max - 30 mg day. |
| HOW TO TAKE | By mouth |
| USES | Bipolar - Anti psychotics, TLE Disorder, Tourette Syndrome |
| SIDE EFFECTS | hypersensitivity - tardive Dyskinesia, stroke |
| WARNINGS | Suicidal - Dementia - Related Psychosis |

| | |
|--------------|--|
| DRUG NAME | Clozaril |
| GENERIC NAME | Clozapine, FazaClo, Versacloz, A |
| DOSAGE RANGE | 12.5 mg increase 100 mg |
| HOW TO TAKE | By mouth |
| USES | Schizophrenia |
| SIDE EFFECTS | Dyslipidemia, Headaches |
| WARNINGS | neutropenia Risk, - Orthostatic Hypotension, SEIZURES, Myocarditis |

| | |
|--------------|---|
| DRUG NAME | Haldo Decanoate |
| GENERIC NAME | Haldo Decanoate, Haldo Lactate → Haloperidol |
| DOSAGE RANGE | Injectable: 0.5mg 1mg 2mg 5mg 10mg, 20mg |
| HOW TO TAKE | Injection. |
| USES | Tourette Syndrome, Agitation, Delirium, anti |
| SIDE EFFECTS | Insomnia, Drowsiness, gynecomastia |
| WARNINGS | Extrapyramidal sx, Leukopenia - Retinopathy |

| | |
|--------------|---|
| DRUG NAME | Dilantin |
| GENERIC NAME | Phenytoin |
| DOSAGE RANGE | ER CAP 30mg, 100mg, Chewable: 50 mg Susp. 125mg per 5mL |
| HOW TO TAKE | By mouth, IV. |
| USES | Seizure Disorders |
| SIDE EFFECTS | Nausea, vomiting, Rash, Ataxia, Headache |
| WARNINGS | Ventricular fib. Hypotension: Stevens-Johnson Synd. |

| | |
|--------------|---|
| DRUG NAME | Levothyroxine |
| GENERIC NAME | Euthyrox, levothyroid, levoxyl, Synthroid, Thyquidity |
| DOSAGE RANGE | 13mg ^{cap} to 300mg Inj. Various. / Hypothyroidism 50-200 mg qd. |
| HOW TO TAKE | By mouth / oral Inj. |
| USES | thyroid Disorders. Hypothyroidism |
| SIDE EFFECTS | Palpitations, Appetite incr. tachycardia tremor, weight loss |
| WARNINGS | not for obesity/weight loss |

| | |
|--------------|---|
| DRUG NAME | Lisinopril |
| GENERIC NAME | Prinivil, Qbrelis, Zestril |
| DOSAGE RANGE | 2.5mg - 40mg |
| HOW TO TAKE | By Mouth |
| USES | Angiotensin-converting (ACE) inhibitors Blood Press |
| SIDE EFFECTS | Anaphylactoid rxn, Stevens-Johnson Syndrome, pancreatitis |
| WARNINGS | Fetal Toxicity |

| | |
|--------------|---|
| DRUG NAME | Flonase |
| GENERIC NAME | Clarispray Anticholinergics, Intranasal Zinc |
| DOSAGE RANGE | 1-2 actuations in Each nostril |
| HOW TO TAKE | 1-2 actuations in Each nostril |
| USES | Corticosteroid for relieving allergic rhinitis symptoms |
| SIDE EFFECTS | Headache, Unise Epistaxis nasal irritation nausea |
| WARNINGS | Hypersensitivity, Anaphylaxis glaucoma, cataracts |

| | |
|--------------|--|
| DRUG NAME | Advair Dises |
| GENERIC NAME | fluticasone propionate |
| DOSAGE RANGE | 100mcg/50mcg - 500mcg/50mcg per Blesster |
| HOW TO TAKE | 1 puff inhaled |
| USES | Asma maintenance |
| SIDE EFFECTS | Bronchospasm, paradoxical, Bothma exacerbation |
| WARNINGS | premenia, laryngospasm |

Black Box

| | |
|--------------|---|
| DRUG NAME | Ativan |
| GENERIC NAME | Lorazepam |
| DOSAGE RANGE | 0.5mg - 4mg per ml |
| HOW TO TAKE | 2-6mg/Day PO By mouth |
| USES | Anxiety, Insomnia, Status Epilepticus |
| SIDE EFFECTS | Sedation, Dizziness, Ataxia, Ataxia - chemo - |
| WARNINGS | Concomitant opioid use. Addiction, Abuse, Misuse. |

| | |
|--------------|--|
| DRUG NAME | Toprol |
| GENERIC NAME | Metoprolol Succinate |
| DOSAGE RANGE | ER TAB 25mg - 200mg |
| HOW TO TAKE | PO By mouth |
| USES | HTN - Angina, Heart Failure, Myocardial infarction |
| SIDE EFFECTS | Fatigue, Dizziness, Diarrhea, pruritis |
| WARNINGS | AVOID Abrupt Cessation |