

Alondra Vidales Lopez



### DMA TRAINING LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [ 48 ]  
See slide 65 in DMA Packette

KM

<b>Mental Illness Anxiety Disorders</b>	<b>Inhalers Allergy / Asthma</b>	<b>Hyperlipidemia Statins</b>	<b>Diabetes Endocrine &amp; Metabolic</b>
Abilify Ativan Clozari Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa	Advair Discus Atrovent Flonase Flovent Loratadine Proventil	Crestor Lipitor Zocor	Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid
<b>Seizures</b>	<b>Gastrointestinal Disorder Constipation</b>	<b>Blood Pressure Meds</b>	<b>Pain &amp; Inflammation</b>
Dilantin Keppra Neurontin Topamax	Colace Miralax Prilosec Protonix Zantac	HCTZ (hydrochlorothiazide) Lisinopril Toprol Tenormin	Flexeril Motrin Norco Tylenol with Codeine Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 2056

DMA Code #2 1818

DMA Code #3 2002

Alondra Vidales Lopez

DRUG NAME	Abilify
GENERIC NAME	Aripiprazole
DOSAGE RANGE	10 to 30 mg/day
HOW TO TAKE	can be taken with or without food. Swallow the regular tablet whole and do not crush, chew, or break the tablet.
USES	Mood disorders, Bipolar disorder, Schizophrenia, Tourette's Syndrome
SIDE EFFECTS	Dizziness, Lightheadedness, Nausea, Vomiting, tiredness
WARNINGS	Thoughts about suicide or dying, panic attacks, trouble sleeping Mood Swings

DRUG NAME	Ativan
GENERIC NAME	Lorazepam
DOSAGE RANGE	0.5mg, 1mg, or 2mg
HOW TO TAKE	By mouth with or without food as directed by your doctor. Taken every day at regular time or as needed.
USES	Anxiety, Sleeplessness, Agitation, Panic Attack
SIDE EFFECTS	Confusion, Aggression, Severe drowsiness, upper stomach pain
WARNINGS	Drowsiness, Blurred Vision, Headache, Nausea

DRUG NAME	Clozaril
GENERIC NAME	Clozapine
DOSAGE RANGE	12.5mg, 25mg, 50mg
HOW TO TAKE	taken 1 or 2 times per day with or without food.
USES	Mood disorders, Schizophrenia, Schizoaffective disorders.
SIDE EFFECTS	drooling, drowsiness, dizziness, Lightheadedness, Headache
WARNINGS	Dry mouth, having trouble keeping your balance

DRUG NAME	Depakote
GENERIC NAME	Divalproex Sodium
DOSAGE RANGE	125mg, 250mg, 500mg
HOW TO TAKE	Swallow Depakote tablets
USES	seizure disorders, bipolar disorder, prevent migraine headaches
SIDE EFFECTS	Vomiting, Abdominal Pain, diarrhea, Low platelet count
WARNINGS	Sleepiness

DRUG NAME	<del>Haloperidol</del> Haldol
GENERIC NAME	Haloperidol
DOSAGE RANGE	50mg, 5mg, 20mg
HOW TO TAKE	Taken 1 or 2 times per day with or without food
USES	treat psychotic disorders like Schizophrenia
SIDE EFFECTS	Dizziness, Dry Mouth, Vomiting, Nausea, Diarrhea
WARNINGS	Lightheadedness, difficulty urinating

Alondra Viales-10/2

DRUG NAME	Invega
GENERIC NAME	paliperidone
DOSAGE RANGE	3mg, 6mg
HOW TO TAKE	Take tablet Swallow the tablet with whole liquid
USES	Schizophrenia, Schizoaffective disorder
SIDE EFFECTS	Upset Stomach, Weight gain, headache, dizziness
WARNINGS	Dizziness and lightheadedness

DRUG NAME	Klonopin
GENERIC NAME	Clonazepam
DOSAGE RANGE	1mg, 2mg
HOW TO TAKE	Swallow tablet without or with food
USES	Seizures and panic disorder
SIDE EFFECTS	Weakness, dizziness, headache, depression
WARNINGS	New or worsening Seizures, Confusion, Aggression

DRUG NAME	Lamictal
GENERIC NAME	Lamotrigine
DOSAGE RANGE	25mg, 50mg, 400mg
HOW TO TAKE	taken 1 or 2 times daily with or without food
USES	epileptic seizures, mood episodes
SIDE EFFECTS	Ataxia, Skin rash, headache, insomnia, Nausea
WARNINGS	Fever and rash, High levels of blood ferritin

DRUG NAME	Lithium
GENERIC NAME	Lithium carbonate
DOSAGE RANGE	20-30mg/Kg
HOW TO TAKE	Swallow tablets with water or juice
USES	prevent Episodes of mania
SIDE EFFECTS	Headaches, Nausea or Vomiting, Diarrhea, Dry mouth
WARNINGS	Change in heart rhythm, muscle weakness, fatigue

DRUG NAME	Risperdal
GENERIC NAME	Risperidone
DOSAGE RANGE	2mg, 3mg, 6mg, 1mg
HOW TO TAKE	take 1 or 2 times per day with or without food Swallow
USES	Bipolar disorder, irritability associated, Autistic disorder
SIDE EFFECTS	Nausea, Vomiting, heartburn, dry mouth, heartburn
WARNINGS	Increased appetite, Increased Saliva