

Shanorria
Walls



DMA TRAINING

LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
See slide 65 in DMA Packette

Mental Illness Anxiety Disorders	Inhalers Allergy / Asthma	Hyperlipidemia Statins	Diabetes Endocrine & Metabolic
Abilify Ativan Clozaril Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa	Advair Discus Atrovent Flonase Flovent Loratadine Proventil	Crestor Lipitor Zocor	Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid
Seizures	Gastrointestinal Disorder Constipation	Blood Pressure Meds	Pain & Inflammation
Dilantin Keppra Neurontin Topamax	Colace Miralax Prilosec Protonix Zantac	HCTZ (hydrochlorothiazide) Lisinopril Toprol Tenormin	Flexeril Motrin Norco Tylenol with Codeine Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 2056

DMA Code #2 1818

DMA Code #3 2002

Shanorria Walls

DRUG NAME	Invega
GENERIC NAME	paliperidone
DOSAGE RANGE	every 6 months between 1,092 mg and 1,560
HOW TO TAKE	by mouth with or without food as directed by your doctor
USES	to treat schizophrenia and schizoaffective disorder
SIDE EFFECTS	abnormal muscle movements
WARNINGS	

DRUG NAME	Klonopin
GENERIC NAME	Clonazepam
DOSAGE RANGE	1.5 mg/day divided into three doses
HOW TO TAKE	May be taken with or without food
USES	Prevent and control seizures
SIDE EFFECTS	depression, headache, weakness,
WARNINGS	

DRUG NAME	Lamictal
GENERIC NAME	Lamotrigine
DOSAGE RANGE	50 mg po qday for 2 weeks then 100 mg/day divided
HOW TO TAKE	by mouth taken w/ or without water or food
USES	mood stabilizer medication treatment of bipolar disorder
SIDE EFFECTS	headache, Diarrhea, Insomnia, Nausea
WARNINGS	

DRUG NAME	Lithium
GENERIC NAME	E Lithicite, Lithobid
DOSAGE RANGE	600 mg 2 times a day or 3 times a day up to 1200 mg
HOW TO TAKE	Swallow tablets whole with a drink of water or juice
USES	To treat the manic episodes of manic depression
SIDE EFFECTS	Headache, Diarrhea, hand tremors, nausea, vomiting
WARNINGS	

DRUG NAME	Risperdal
GENERIC NAME	Risperidone
DOSAGE RANGE	2mg per day
HOW TO TAKE	It can treat schizophrenia, bipolar, and irritability
USES	Take by mouth once or twice a day
SIDE EFFECTS	Dry mouth, heart burn, nausea, Diarrhea, vomiting
WARNINGS	

Shanorria
Wall

DRUG NAME	Abilify
GENERIC NAME	Aripiprazole
DOSAGE RANGE	2 to 5 mg/day. Dosage adjustments up to 5mg/day
HOW TO TAKE	
USES	take medication by mouth ^{with} or without food as
SIDE EFFECTS	dizziness, lightheadedness, drowsiness, nausea, vomiting
WARNINGS	compulsive or uncontrollable urges to gamble, binge, eat, shop and have sex

DRUG NAME	Ativan Ativan
GENERIC NAME	Lorazepam
DOSAGE RANGE	2 to 6 mg/day given in divided doses
HOW TO TAKE	may be taken with or without food blurred vision
USES	
SIDE EFFECTS	drowsiness, dizziness, loss of coordination Headache, nausea
WARNINGS	

DRUG NAME	Clozaril
GENERIC NAME	Clozapine
DOSAGE RANGE	12.5 mg once daily or twice daily
HOW TO TAKE	take by mouth once or twice daily
USES	
SIDE EFFECTS	
WARNINGS	drowsiness dry mouth headache restless increased salivation

DRUG NAME	Depakote
GENERIC NAME	Divalproex sodium
DOSAGE RANGE	250 mg twice daily
HOW TO TAKE	Swallow whole or sprinkle onto food
USES	
SIDE EFFECTS	
WARNINGS	fever, flu symptoms, runny nose, sore throat

DRUG NAME	Haldol
GENERIC NAME	Haloperidol
DOSAGE RANGE	0.5-2 mg q8-12hr initially
HOW TO TAKE	take 1 or 2 times per day with or without food
USES	
SIDE EFFECTS	dry mouth, blurred vision, diarrhea, heartburn
WARNINGS	