



## Medication Administration In-Service and Evaluation

Name of Facility/Home: Anchor Point

Employee Receiving In-Service: Kailie Sargent

Date of 1st In-Service: 8 / 19 / 202 Time: 1 : 00pm am / pm Trainer: Learning and Development

Date of 2nd In-Service: 8 / 19 / 202 Time: 1 : 00pm am / pm Trainer: Learning and Development

Date of 3rd In-Service: 9 / 21 / 21 Time: 2 : 00 am / (pm) Trainer: Flora Lawton

Date of 4th In-Service: 9 / 30 / 21 Time: 3 : 00 am / (pm) Trainer: Kelly Greer

Date of 5th In-Service: 9 / 30 / 21 Time: 3 : 30 am / (pm) Trainer: Kelly Greer

Date of 6th In-Service: 10 / 01 / 21 Time: 4 : 00 am / (pm) Trainer: Flora Lawton

Date of Final Evaluation: 10 / 01 / 21 Time: 8 : 00 am (pm) Trainer: Flora Lawton

**All staff must complete all three (6) In-Services and Final Evaluation**

Instructions: Check (✓) the appropriate box after Employee has been in-serviced.

		In-Service #	1st	2nd	3rd	4th	5th	6th	Eval.	Comments
1	Medication Area		<input checked="" type="checkbox"/>							
	a. Location of ample supplies prior to administration		<input checked="" type="checkbox"/>							
	b. Area is clean and organized		<input checked="" type="checkbox"/>							
	c. Area is always locked		<input checked="" type="checkbox"/>							
	d. Location of all medication: Internal, External, Refrigerated, Controlled Drugs (narcotics)		<input checked="" type="checkbox"/>							
2	DMA washes hands prior to administering medications and between each Resident		<input checked="" type="checkbox"/>							
3	Medication keys are retained by DMA		<input checked="" type="checkbox"/>							
4	Resident is identified per facility policy and procedure prior		<input checked="" type="checkbox"/>							
5	Vital signs are taken per facility policy prior to administering medications (if applicable), always on cardiac and BP medications		<input checked="" type="checkbox"/>							
	a. If Pulse and BP are required, hands and equipment are washed per facility policy		<input checked="" type="checkbox"/>							
	b. If Apical Pulse is required, privacy is provided		<input checked="" type="checkbox"/>							
6	Medications Administration per facility policy and procedure: to include review of the '6 Rights'		<input checked="" type="checkbox"/>							
	a. Medications are properly removed from container/blister pack and (.) dot is placed in appropriate box on MAR		<input checked="" type="checkbox"/>							
	b. Liquid medication is poured at eye level, with palm covering label of stock bottle		<input checked="" type="checkbox"/>							



## Medication Administration In-Service and Evaluation

In-Service #	Description	1st	2nd	3rd	4th	5th	6th	Eval.	Comments
6	c. DMA verifies medication and strength with order as transcribed on medication record per facility policy and procedure	<input checked="" type="checkbox"/>							
	d. Observe Resident to ensure medication is swallowed	<input checked="" type="checkbox"/>							
	e. Offer adequate and appropriate fluid with medication	<input checked="" type="checkbox"/>							
	f. Medication record is signed immediately after administration of same	<input checked="" type="checkbox"/>							
	g. Controlled substance record is signed immediately after administration of same	<input checked="" type="checkbox"/>							
	h. Correct dose is administered	<input checked="" type="checkbox"/>							
	i. Medication is administered at correct time	<input checked="" type="checkbox"/>							
	j. Verify no additional MAR pages have been added	<input checked="" type="checkbox"/>							
7	Infection control technique is reviewed	<input checked="" type="checkbox"/>							
8	Medication via gastric tube administered per facility policy and procedure (if applicable)	<input checked="" type="checkbox"/>							
	a. Resident is properly positioned, at a 45° sitting angle	<input checked="" type="checkbox"/>							
	b. Tube is checked for placement and patency	<input checked="" type="checkbox"/>							
	c. Tube is flushed before, between and after medications are administered	<input checked="" type="checkbox"/>							
9	Injections are administered by the Resident or a DMA if there is a doctor's order present, per facility policy and procedure	<input checked="" type="checkbox"/>							
	a. Syringes and needles are disposed of in sharps container, by person giving the injection without recapping	<input checked="" type="checkbox"/>							
	b. Proper glucometer testing is observed. Determination of competence re: accurately perform and read glucometer testing results	<input checked="" type="checkbox"/>							
10	DMA crushes medication according to facility policy and procedure ONLY with physician's orders.	<input checked="" type="checkbox"/>							
11	DMA administers eye and ear medication according to facility policies and procedures	<input checked="" type="checkbox"/>							
12	Side effects of psychoactive medication are noted (lethargy, hallucinations) and reported.	<input checked="" type="checkbox"/>							
13	Medication administration should not interrupted. DO NOT RUSH	<input checked="" type="checkbox"/>							
14	Controlled drugs are stored (Double Locked) according to facility policy and procedure	<input checked="" type="checkbox"/>							
15	Residents' rights are observed	<input checked="" type="checkbox"/>							
16	Location, Procedures and Documenting for administering PRN	<input checked="" type="checkbox"/>							
17	Designated Medication Administrator follows facility policy and procedure for medications refused or withheld. (MER & IR written)	<input checked="" type="checkbox"/>							
18	Medications are administered within time frame per facility policy	<input checked="" type="checkbox"/>							



## Medication Administration In-Service and Evaluation

	In-Service #	1st	2nd	3rd	4th	5th	6th	Eval.	Comments
19	Medication errors are reported to Site Supervisor and RN teaching medication classes	<input checked="" type="checkbox"/>							
20	Medication area is cleaned and locked after completion of medication administration	<input checked="" type="checkbox"/>							
21	Designated Medication Administrator can identify action and common side effects of medications administered	<input checked="" type="checkbox"/>							
22	Approved Abbreviations List is reviewed	<input checked="" type="checkbox"/>							
23	Seizure precautions and documentation	<input checked="" type="checkbox"/>							
24	After hour procedures, procedures for found/spilled medication, location of Guide to Drugs Book	<input checked="" type="checkbox"/>							
25	2nd Staff Verification, what it is, when it is needed, and how to document it	<input checked="" type="checkbox"/>							
26	Refusal of Medication procedures (prompt 3 times, then write appropriate documentation)	<input checked="" type="checkbox"/>							

### FOLLOW UP CONCERNS

Specify time frame for completion: \_\_\_\_\_  N/A

---



---



---

I have received the above In-service and have read the Organizations **Medical Policies**. I understand what is expected of me as a Designated Medication Administrator. I also understand that any immediate medical questions or concerns should be directed to the Coordinator of Care at my Site during open office hours and to the On-Call person after hours.

Kailie Sargent  
Employee Signature

10-1-21  
Date

[Signature]  
Home Manager Signature

10-1-21  
Date

## How to Register for an Epocrates Account

**Objective:** Register for Epocrates.com

**User Security Access Requirements:** Available to all staff


**Beacon User Role:** Available to all staff

**Timeline:** Should create an account upon initiating DMA certification training process.


### How to Register:

Pull up Google Chrome or any other web browser and go to <https://www.epocrates.com/>

You will now be the Epocrates home page.



Click on "Login" in the upper right-hand corner of the page.



DRUG NAME	Klonopin
GENERIC NAME	Clonazepam
DOSAGE RANGE	<del>0.5mg, 1mg, 2mg</del> 0.25mg - 2mg
HOW TO TAKE	Start 0.25mg PO bid, incr. by 0.25mg/day q 1-2 days
USES	max: 4mg/day - lower start dose in elderly pts taper grad.
SIDE EFFECTS	drowsiness, dizziness, depression, fatigue, impaired coordination
WARNINGS	abuse and misuse can lead to overdose and death

DRUG NAME	Flovent (HFA)
GENERIC NAME	Fluticasone propionate inhaled
DOSAGE RANGE	2-4 puffs inhaled
HOW TO TAKE	start 2 puffs inhaled b.i.d, may incr dose after 2 wks
USES	to prevent asthma attacks in adults and children
SIDE EFFECTS	URI sx, headache, throat irritation, candidiasis (oral)
WARNINGS	Do not use if allergic to milk proteins

DRUG NAME	Keppra
GENERIC NAME	levetiracetam
DOSAGE RANGE	500 - 1500 mg
HOW TO TAKE	Start 500mg PO/IV q 12h, may incr. by 1000mg/day
USES	treat certain types of seizures in people with epilepsy
SIDE EFFECTS	Some people have thoughts about suicide
WARNINGS	Seizures may increase if you stop taking sudden

DRUG NAME	Prilosec
GENERIC NAME	Omeprazole
DOSAGE RANGE	10 - 40mg
HOW TO TAKE	give or take before meals
USES	to treat symptoms of gastroesophageal reflux (GERD)
SIDE EFFECTS	may cause new or worsening symptoms of lupus
WARNINGS	stop using if you have stomach pain, diarrhea new or unusual pain

DRUG NAME	NORCO
GENERIC NAME	hydrocodone / acetaminophen
DOSAGE RANGE	2.5 - 10 mg
HOW TO TAKE	Follow all directions on your prescription label
USES	Pain, moderate - mod. severe
SIDE EFFECTS	dizziness, nausea, constipation, headache, stomach
WARNINGS	Taking opioid medication during pregnancy may cause i.e. thrombolytic withdrawal symptoms in newborns

DRUG NAME	To pa max
GENERIC NAME	Topiramate
DOSAGE RANGE	15 - 200 mg
HOW TO TAKE	Swallow tablet whole and do not crush, chew, or break
USES	used to treat certain types of seizures in adults
SIDE EFFECTS	can make birth control pills less effective
WARNINGS	can increase body temperature and decrease sweating which may lead to life threatening dehydration

DRUG NAME	Zocor
GENERIC NAME	Simvastatin
DOSAGE RANGE	5 - 80 mg
HOW TO TAKE	take <del>oral</del> oral suspension on an empty stomach
USES	to lower blood levels with bad cholesterol
SIDE EFFECTS	muscle weakness, trouble lifting arms, liver problems
WARNINGS	do not take if pregnant or breastfeeding

DRUG NAME	Lisinopril
GENERIC NAME	Lisinopril
DOSAGE RANGE	2.5 - 40 mg
HOW TO TAKE	measure liquid medicine carefully using measuring device
USES	used to treat high blood pressure
SIDE EFFECTS	Fever, kidney and liver problems, light headed
WARNINGS	do not take within 36hrs before or after taking medication that contains sacubitril

DRUG NAME	Lantus
GENERIC NAME	insulin glargine
DOSAGE RANGE	100 units per ml
HOW TO TAKE	Injected under the skin
USES	to improve blood sugar control in people with <sup>diabetes</sup>
SIDE EFFECTS	rapid weight gain, shortness of breath, low potassium
WARNINGS	must not be given with an insulin pump

DRUG NAME	Novolog
GENERIC NAME	insulin aspart
DOSAGE RANGE	100 units per ml
HOW TO TAKE	injected under skin or infusion into a vein
USES	a hormone that works by lowering glucose levels
SIDE EFFECTS	heart problems, swelling, skin rash
WARNINGS	not approved for use by anyone younger than 2yrs