

Ihuassi Badet

DRUG NAME	ABILIFY
GENERIC NAME	ARIPIPRAZOLE
DOSAGE RANGE	10-15mg/day
HOW TO TAKE	Can be taken with or without food and swallow it
USES	It uses to treat schizophrenia
SIDE EFFECTS	dystonia, stroke, syncope, seizures ---
WARNINGS	nausea, drowsiness, vomiting, dizziness.

DRUG NAME	CLOZARIL
GENERIC NAME	CLOZAPINE
DOSAGE RANGE	25-50mg/day
HOW TO TAKE	It is taken orally in tablet form
USES	It is used to treat schizophrenia
SIDE EFFECTS	Constipation, drooling, shaking, weight gain ---
WARNINGS	Myocarditis, severe neutropenia ---

DRUG NAME	DEPAKOTE
GENERIC NAME	DIVALPROEX SODIUM
DOSAGE RANGE	250mg twice daily
HOW TO TAKE	Swallow it, take it with food if stomach upset occurs
USES	Used to treat seizure disorders, bipolar disorder ---
SIDE EFFECTS	Diarrhea, dizziness, hair loss, shakiness ---
WARNINGS	Hepatic failure, pancreatitis, Hypothermia ---

DRUG NAME	HALDOL
GENERIC NAME	HALOPERIDOL
DOSAGE RANGE	2-5mg, maximum 20mg/day
HOW TO TAKE	Swallow it
USES	Used to treat schizophrenia and psychosis
SIDE EFFECTS	Agitation, hallucinations, fever, chills ---
WARNINGS	Cardiovascular effects, falls, hypersensitivity reactions

DRUG NAME	INVEGA
GENERIC NAME	PALIPERIDONE
DOSAGE RANGE	6-12mg/day
HOW TO TAKE	by mouth with or without food
USES	It is used to treat Schizoph
SIDE EFFECTS	drooling, stomach pain, tiredness, dizziness, ---
WARNINGS	tremors, weight gain, anxiety ---

DRUG NAME	KLONOPIN
GENERIC NAME	CLONAZEPAM
DOSAGE RANGE	0.5 - 1mg / day
HOW TO TAKE	Administered with water by swallowing the tablet
USES	Used to treat the symptoms of seizure and panic disorder
SIDE EFFECTS	Confusion, hallucinations, weak breathing ---
WARNINGS	Depression, porphyria, sleep apnea

DRUG NAME	LORATADINE
GENERIC NAME	Alavert
DOSAGE RANGE	10 mg / day
HOW TO TAKE	Swallow it.
USES	Used to treat sneezing, hives, skin rash, itching.
SIDE EFFECTS	headache, drowsy, dry mouth, hyperactive
WARNINGS	Asthma, liver disease, kidney disease ---

DRUG NAME	CRESTOR
GENERIC NAME	ROSUVASTATIN
DOSAGE RANGE	10 - 20 mg
HOW TO TAKE	with water by swallowing.
USES	Used to reduce or lower cholesterol
SIDE EFFECTS	depression, headache, diarrhea, nausea, joint pain
WARNINGS	hypothyroidism, myopathy, alcohol abuse

DRUG NAME	LANTUS
GENERIC NAME	INSULIN GLARGINE
DOSAGE RANGE	0, 2 units / kg / dose
HOW TO TAKE	Injection
USES	Used to treat diabetes
SIDE EFFECTS	hypoglycemia, headache, hunger, tremors ---
WARNINGS	renal impairment, hypokalemia, hypoglycemia

DRUG NAME	NORCO
GENERIC NAME	HYDROCODONE
DOSAGE RANGE	1 or 2 tablets every four to six hours as needed
HOW TO TAKE	With water by swallowing.
USES	Used to relieve relieve severe pain
SIDE EFFECTS	Noisy breathing, Sighing, nausea, tiredness
WARNINGS	Seizure disorder, head injury, circulatory shock.