

DRUG NAME	Ativan	Diane Harris
GENERIC NAME	Lorazepam	
DOSAGE RANGE	0.5 mg, 1 mg, 2 mg (injection IV) 4 mg per ml	
HOW TO TAKE	with or without food, every day at regular times or on an as needed basis.	
USES	Anxiety disorder and panic attacks for sleep/insomnia and seizures	
SIDE EFFECTS	Somnolence, dizziness, loss of coordination	
WARNINGS	Change in sexual interest/libido, constipation, heartburn, or changes in appetite may occur.	

DRUG NAME	Halodol
GENERIC NAME	Haloperidol
DOSAGE RANGE	0.5 mg, 1 mg, 2 mg, 5 mg, 10 mg and 20 mg
HOW TO TAKE	Taken 1 or 2 times per day with or without food
USES	A psychiatric medication that works by helping to restore the balance in the brain
SIDE EFFECTS	Mood changes, agitation, hallucinations, drowsiness, trouble breathing or swallowing
WARNINGS	Increased mortality in elderly patients with dementia

DRUG NAME	Seroquel
GENERIC NAME	Quetiapine
DOSAGE RANGE	400-800 mg a day / or 600 mg a day / 25 mg
HOW TO TAKE	Usually 1, 2, or 3 times a day
USES	For bipolar disorder / anxiety
SIDE EFFECTS	Dry mouth
WARNINGS	There may be a slightly increased risk of serious, possibly fatal side effects

DRUG NAME	Atraxent
GENERIC NAME	Ipratropium bromide
DOSAGE RANGE	17 mg (actuation) / 0.03% nasal spray / 0.02% nebulized solution
HOW TO TAKE	Prescribed by doctor
USES	Used to treat and prevent symptoms of wheezing and shortness of breath.
SIDE EFFECTS	Headaches, dry mouth, hoarseness, cough, stuffy nose, sinus pain, nausea, upset
WARNINGS	Do not get in your eyes.

DRUG NAME	Celacephage
GENERIC NAME	Metformin
DOSAGE RANGE	500 mg, 850 mg, 1000 mg, 1500 mg, 2550 mg
HOW TO TAKE	Take with a meal, unless your doctor says otherwise
USES	It can treat type 2 diabetes
SIDE EFFECTS	Heartburn, nausea or vomiting, bloating, stomach pain
WARNINGS	Lactic acidosis is a rare, but serious side effect.

DRUG NAME	Zilexon
GENERIC NAME	Cyclobenzaprine
DOSAGE RANGE	5 mg, 10 mg, 15 mg 30 mg
HOW TO TAKE	
USES	is a muscle relaxant.
SIDE EFFECTS	Dry mouth, heartburn, extreme tiredness, dizziness
WARNINGS	Do not take this medication if you have an allergy to the medication.

DRUG NAME	Drilosec
GENERIC NAME	Omeprazole
DOSAGE RANGE	20 mg
HOW TO TAKE	Used to treat certain stomach & esophagus problems
USES	Take this medication by mouth
SIDE EFFECTS	Headaches, nausea, vomiting, stomach pain, gas, fever, etc.
WARNINGS	Your risk may be higher if you take this medication.

DRUG NAME	Sifantin
GENERIC NAME	Phenytoin
DOSAGE RANGE	30 mg, 100 mg (capsules) or pediatric 5 mg / 4-8 mg
HOW TO TAKE	Take this medication by mouth usually once a day or in divided doses as directed
USES	Prevent and control seizures
SIDE EFFECTS	Headaches, nausea, vomiting, constipation, feeling of spinning
WARNINGS	Do not take if you take delavirdine (Rescriptor) or if you are allergic to phenytoin, ethoin, fosphenytoin, ethoin or if you are pregnant

DRUG NAME	Lipitor
GENERIC NAME	Atorvastatin
DOSAGE RANGE	10-80 mg daily
HOW TO TAKE	orally once daily
USES	used for the treatment of elevated total cholesterol, LDL, and to elevate HDL cholesterol
SIDE EFFECTS	constipation, diarrhea, nausea, gas, fatigue, heartburn
WARNINGS	Do not take if you are allergic to atorvastatin or any of the ingredients contain in this drug

DRUG NAME	Toprol
GENERIC NAME	Metoprolol
DOSAGE RANGE	50 mg orally twice a day / 100 mg orally once a day
HOW TO TAKE	you'll take by mouth
USES	To treat high blood pressure, chest pain (angina), and heart failure.
SIDE EFFECTS	dizziness or light headedness, tiredness, dryness
WARNINGS	you should not use this medication if you have a serious heart problem, severe heart beats that caused fainting, severe circulation problems