

DMA Code 1: 0978

DMA Code 2: 0439

DMA Code 3: 0501

a) Drug Name	Abilify	g) How to take: PO / by mouth qd
b) Generic Name	aripiprazole	
c) Dosage Range	2-30 mg	
d) Uses	Schizophrenia, bipolar I, major depressive disorder	
e) Side Effects	Hyperglycemia, cough, fatigue, diarrhea, insomnia, weight gain, etc.	
f) Warnings	Not approved for dementia related psychosis, increased suicidality	

2. a) Haldol
 b) haloperidol
 c) 0.5-20 mg
 d) psychosis, Tourette's, agitation/acute, renal & hepatic dosing
 e) Hypotension, insomnia, anxiety, lethargy, photosensitivity, menstrual irregularities, etc.
 f) not approved for dementia related psychosis
 g) PO bid-tid

3. a) Flonase Allergy Relief
 b) fluticasone propionate nasal
 c) allergy: 1-2 actuations each nostril qd/day; nasal polyp tx: 2-4 actuations each nostril ^{bid/day}
 d) allergy, nasal polyps
 e) Hypersensitivity rxn, anaphylaxis, headache, nasal burning or irritation, bronchitis, etc.
 f) If patient has unhealed nasal septal ulcer or wound, long-term pediatric use, hepatic impairment, etc.
 g) Nostril bid/qd ~~(strength)~~ actuations

4. a) Lipitor
 b) atorvastatin
 c) 10-80 mg
 d) Hypercholesterolemia, mixed dys-lipidemia, hypertriglyceridemia type IV, dysbetalipoproteinemia etc.
 e) Myopathy, tendon rupture, URI, UTI, Nausea, insomnia, muscle spasms, myalgia, etc.
 f) ~~Not for~~ If patient has hypothyroidism, stroke w/in 6mo, diabetes mellitus, etc.
 g) PO qd

5. a) Synthroid
 b) Levothyroxine
 c) 25-300 mcg
 d) Hypothyroidism, TSH suppression
 e) Hypersensitivity, arrhythmia, palpitations, tachycardia, weight loss, tremor, etc.
 f) Not for obesity/weight loss, caution if pregnant or hyperactivity risk, TSH suppression, etc.
 g) PO qd

6. a) Keppra
 b) levetiracetam
 c) 250-1000 mg, SOL: 100 mg/mL
 d) partial seizures, juvenile myoclonic epilepsy, primary generalized tonic clonic seizures
 e) Depression, hostility, vomiting, headache, asthenia, fatigue, anorexia, psychosis, etc.
 f) Avoid abrupt withdrawal, caution if pregnant, if CrCl < 80, if psychiatric disorder present etc
 g) PO/IV q12h

7. a) Gluco phage
 b) metformin
 c) 500 mg - 1000 mg
 d) diabetes mellitus / type 2, polycystic ovary syndrome
 e) Lactic acidosis, sepsis, surgery, caution if premenopausal anovulatory pts, etc.
 f) Megaloblastic anemia, diarrhea, nausea, vomiting, anorexia, rash, headache, etc.
 g) PO qd/bid/tid

8. a) Colace
 b) docusate sodium
 c) 2-11 yo: 50-150 mg/day, 12 yo ↑: 50-300 mg/day
 d) Constipation, stool softener, emollient laxative (PO liquid forms)
 e) Diarrhea, abdominal cramps, throat irritation, rash, electrolyte disorders, bitter taste
 f) Fecal impaction, mineral oil use, appendicitis, acute abdomen, GI obstruction, etc.
 g) PO prn

9. a) Toprol
 b) metoprolol succinate
 c) 25-200 mg
 d) HTN, angina, heart failure w/ reduced ejection fraction / NYHA class II-IV, migraine headache prophylaxis
 e) Severe bradycardia, fatigue, pruritus, dyspnea, depression, nightmares, etc.
 f) Avoid abrupt cessation, HR < 45 bpm, 2nd/3rd degree AV block, cardiogenic shock, etc.
 g) PO qd

10. a) ~~Flexril~~ ← (discontinued in US market, interesting) instead → Motrin
 b) ~~cyclobenzaprine~~ ibuprofen
 c) 1-2 tabs/caps q4-6h prn, max 6 tabs/caps/24h (use lowest effective dose)
 d) NSAID for reducing pain/inflammation, fever
 e) GI bleeding, perforation, or ulcer; stroke, nausea, rash, tinnitus, constipation, etc.
 f) CrCl < 30, Aspirin tirad; caution if pregnant, in elderly pts, debilitated pts, etc.
 g) PO q4-6h prn