

Saralee Noble

## DMA TRAINING

### LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [ 48 ]  
See slide 65 in DMA Packette

Mental Illness Anxiety Disorders	Inhalers Allergy / Asthma	Hyperlipidemia Statins	Diabetes Endocrine & Metabolic
Abilify Ativan ✓ Clozaril Depakote ✓ Haldol Invega Klonopin Lamictal ✓ Lithium Risperdal Seroquel Tripleptal Zyprexa	Advair Discus ✓ Atrovent Flonase Flovent Loratadine Proventil	Crestor ✓ Lipitor Zocor	Apidra Byetta Glucophage Glyburide ✓ Lantus Levemir Levothyroxine ✓ Novolog Synthroid
Seizures	Gastrointestinal Disorder Constipation	Blood Pressure Meds	Pain & Inflammation
Dilantin Keppra ✓ Neurontin Topamax	Colace Miralax ✓ Prilosec Protonix Zantac	HCTZ (hydorchlorothiazide) ✓ Lisinopril Toprol Tenormin	Flexeril ✓ Motrin Norco Tylenol with Codeine ✗ Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 0917

DMA Code #2 1017

DMA Code #3 1109

DRUG NAME	<del>clozapil</del> Clozaril
GENERIC NAME	Clozapine, FazaClo, Versacloz
DOSAGE RANGE	Tab - 25mg, 100mg
HOW TO TAKE	Start 12.5mg, increase by 25-50mg/day to target 300-450mg/day in divided doses by day 14, then may incr. by upto 100mg/day
USES	Schizophrenia, tx resistant,
SIDE EFFECTS	Severe Neutropenia, Seizures, Mitral Valve Incompleteness, <sup>Dementia,</sup>
WARNINGS	restricted distribution program due to severe neutropenia risk.

DRUG NAME	Lithium
GENERIC NAME	eskalith, Lithobid, lithium SOL: 8mEq per 5ml
DOSAGE RANGE	Cap: 150mg, 300mg, 600mg, Tab: 300mg, ER Tab: 300mg <sup>450mg</sup>
HOW TO TAKE	See back ->
USES	Bipolar disorder
SIDE EFFECTS	coma, seizures, ventricular arrhythmia, bradycardia, syncope, Brugada Syndrome, drug rxn, goiter, hypothyroidism
WARNINGS	lithium toxicity closely related to serum lithium levels and can occur at doses close to therapeutic levels.

DRUG NAME	Atrovent
GENERIC NAME	Clozapine, revefenacin inhaled
DOSAGE RANGE	<del>MDI</del> MDI: 17mcg per actuation 2 puffs inhaled, <sup>max</sup> 12 puffs/day
HOW TO TAKE	inhale puff
USES	asthma
SIDE EFFECTS	hypersensitivity rxn, anaphylaxis, bronchospasm, paradoxical, glaucoma, angle-closure.
WARNINGS	N/A

DRUG NAME	Lipitor
GENERIC NAME	atorvastatin
DOSAGE RANGE	Tab: 10mg, 20mg, 40mg, 80mg
HOW TO TAKE	Tab form, depending on uses.
USES	hypercholesterolemia, mixed dyslipidemia, hypertriglyceridemia, dysbetalipoproteinemia
SIDE EFFECTS	myopathy, tendon rupture, rhabdomyolysis, acute renal failure, hepatotoxicity, pancreatitis, anaphylaxis, photosensitivity
WARNINGS	N/A

DRUG NAME	Lantus
GENERIC NAME	Insulin glargine
DOSAGE RANGE	100 units per ml
HOW TO TAKE	Injection
USES	diabetes
SIDE EFFECTS	hypersensitivity rxn, anaphylaxis, hypoglycemia, hypokalemia
WARNINGS	N/A

DRUG NAME	Novolog
GENERIC NAME	Fiasp, Novolog, Insulin aspart.
DOSAGE RANGE	100 units per ml
HOW TO TAKE	0.5-1 - total daily insulin requirement give 5-10 mins before meals (sc injections) or continuous infusion (sc pump)
USES	<del>diabetes</del> diabetes
SIDE EFFECTS	hypoglycemia, hypokalemia, hypersensitivity rxn, anaphylaxis.
WARNINGS	N/A

DRUG NAME	Neurontin
GENERIC NAME	gabapentin
DOSAGE RANGE	Cap: 100mg, 300mg, 400mg, Tab: 600mg, 800mg, Sol: 50mg per ml
HOW TO TAKE	cap, tab, sol - Depending on usage.
USES	partial seizures, post-herpetic neuralgia, neuropathic pain, fibromyalgia, alcohol dependence.
SIDE EFFECTS	depression, suicidality, erythema multiforme, Stevens-Johnson syndrome, anaphylaxis, angioedema, rhabdomyolysis, respiratory depression
WARNINGS	N/A

DRUG NAME	PriLOSEC
GENERIC NAME	omeprazole
DOSAGE RANGE	DR CAP: 10mg, 20mg, 40mg, DR GRANULE: 10mg per phl
HOW TO TAKE	give before meals; See PPI Administration Alternatives table
USES	GERD, H. pylori infection, gastric ulcer prophylaxis, <del>gastric ulcer</del>
SIDE EFFECTS	hypersensitivity rxn, anaphylaxis, Stevens-Johnson syndrome, toxic epidermal, erythema multiforme, tubulointerstitial nephritis, renal papillary necrosis
WARNINGS	No Black Box Warnings!

DRUG NAME	Lisinopril
GENERIC NAME	Prinivil, Qbrelis, Zestril, lisinopril
DOSAGE RANGE	TAB: 2.5mg, 5mg, 10mg, 20mg, 30mg, 40mg
HOW TO TAKE	Take tab dosage Depending on Usage.
USES	HTN, Heart failure w/reduced ejection fraction, MI, acute, migraine headache prophylaxis.
SIDE EFFECTS	anaphylactoid rxn, angioe-head or neck-intestinal, Stevens-Johnson syndrome, toxic epidermal necrolysis, hypotension, severe hyperkalemia.
WARNINGS	Fetal/neonatal morbidity/mortality may occur when drugs that act directly on the renin-angiotensin system are used in pregnancy

DRUG NAME	Ultram-discontinued (Motrin)
GENERIC NAME	Ibuprofen
DOSAGE RANGE	12 years or older - (oral tablet/capsule)
HOW TO TAKE	<del>reducing pain/inflammation</del> 1-2 tab/caps Max: 6 tabs/caps 24hrs.
USES	reducing pain/inflammation/fever
SIDE EFFECTS	GI bleeding, GI perforation, GI ulcer, MI, stroke, thromboembolism, HTN, CHF, renal papillary necrosis, nephrotoxicity, hepatotoxicity
WARNINGS	N/A