Darrell Stancie l

DMA TRAINING
LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
See slide 65 in DMA Packette

Mental Illness **Anxiety Disorders**	**Inhalers** **Allergy / Asthma**	**Hyperlipidemia** **Statins**	**Diabetes** **Endocrine & Metabolic**
Abilify Ativan Clozaril Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa	Advair Discus Atrovent Flonase Flovent Loratadine Proventil	Crestor Lipitor Zocor	Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid
Seizures	**Gastrointestinal Disorder** **Constipation**	**Blood Pressure Meds**	**Pain & Inflammation**
Dilantin Keppra Neurontin Topamax	Colace Miralax Prilosec Protonix Zantac	HCTZ (hydorchlorothiazide) Lisinopril Toprol Tenormin	Flexeril Motrin Norco Tylenol with Codeine Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 9574 DMA Code #2 5238 DMA Code #3 1763

DRUG NAME	Ativan
GENERIC NAME	lorazepam
DOSAGE RANGE	Tab 0.5mg, 1mg, 2mg; INJ: 2mg per ML, 4mg per ML
HOW TO TAKE	By mouth
USES	Anxiety, insomnia, status Epilepticus, Nause/vomiting
SIDE EFFECTS	Respiratory depression, apnea, Respiratory failure, abuse, seizures
WARNINGS	avoid use of IM/IV forms in neonates, serious potential fatal gasping syndromes may occur

DRUG NAME	Clozaril
GENERIC NAME	clozapine
DOSAGE RANGE	Tab 25 mg, 100 mg
HOW TO TAKE	By mouth
USES	Schizophrenia, suicide prevention, schizophrenia-assoc
SIDE EFFECTS	agranulocytosis, neutropenia, seizures, myocarditis, cardiomyopathy
WARNINGS	severe neutropenia, seizures

DRUG NAME	Topamax
GENERIC NAME	topiramate
DOSAGE RANGE	Sprinkle cap 15mg, 25mg, Tab: 25mg, 50mg, 100mg, 200mg
HOW TO TAKE	By mouth
USES	partial seizures, seizure primary generalized tonic clonic
SIDE EFFECTS	metabolic acidosis, severe, nephrolithiasis, osteomalacia, osteoporosis
WARNINGS	depression, Behavior changes, suicidality

DRUG NAME	Lipitor
GENERIC NAME	atorvastatin
DOSAGE RANGE	Tab 10mg, 20mg, 40mg, 80mg
HOW TO TAKE	By mouth
USES	hypercholesterolemia, mixed dyslipidemia, hypertriglycer
SIDE EFFECTS	Myopathy, tendon rupture, Rhabdomyolysis, acute renal failure
WARNINGS	CK at baseline if Myopathy Risk

DRUG NAME	Crestor
GENERIC NAME	Rosuvastatin
DOSAGE RANGE	Tab 5mg, 10mg, 20mg, 40mg
HOW TO TAKE	By mouth
USES	Hypercholesterolemia, hypertriglyceridemia, type IV, mixed dyslipid
SIDE EFFECTS	Myopathy, pancreatitis, acute renal, tendon Rupture, hepatotoxicity
WARNINGS	CK at basline if myopath Risk

DRUG NAME	Dilantin
GENERIC NAME	phenytoin
DOSAGE RANGE	ER cap; 30mg, 100mg, chewable; 50mg, Susp; 125mg/5mL
HOW TO TAKE	By mouth
USES	Status epilepticus, seizure disorder
SIDE EFFECTS	ventricular fibrillation, hypotension, severe, cardiac arrest, bradycardia
WARNINGS	depression, behavior, changes, suicidality

DRUG NAME	Colace
GENERIC NAME	docusate sodium
DOSAGE RANGE	2-11 yo, 50-150 mg/day po divided qd bid pm, 12yo, 50-300
HOW TO TAKE	By mouth
USES	Renal dosing, Hepatic dosing
SIDE EFFECTS	diarrhea, abdominal cramps, throat irritation, Rash, Electrolyte disorder
WARNINGS	none Reported

DRUG NAME	Toprol-XL
GENERIC NAME	metoprolol succinate
DOSAGE RANGE	ER Tab: 25mg, 50mg, 100mg, 200mg
HOW TO TAKE	By mouth
USES	angina, heart failure w/reduce ejection fraction
SIDE EFFECTS	Bradycardia, severe, CHF, heart block, cardiogenic shock (misuse)
WARNINGS	Avoid Abrupt cessation

DRUG NAME	NOVOLog
GENERIC NAME	insulin aspart
DOSAGE RANGE	INJ (penfill U-100 cartridge): 100 units per mL, INJ (Flexpen U-100 pen
HOW TO TAKE	By mouth
USES	diabetes mellitus, renal dosing
SIDE EFFECTS	Hypoglycemia, Hypokalemia, Hypersensitivity RXN, anaphlaxis
WARNINGS	caution in pts 65yo, w/o concurrent basal or long acting insulin

DRUG NAME	Keppra
GENERIC NAME	levetiracetam
DOSAGE RANGE	TAB: 250mg, 500mg, 750mg, 1000mg, SOL: 100mg per mL
HOW TO TAKE	By mouth
USES	partial seizures, Juvenile myoclonic epilepsy, adjunct tx
SIDE EFFECTS	depression, behavior changes, Suicidality, hostility, psychosis
WARNINGS	depression, Sucidality,