

Melissa

DMA TRAINING
LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48] See slide 65 in DMA Packette

Mental Illness Anxiety Disorders	Inhalers Allergy / Asthma	Hyperlipidemia Statins	Diabetes Endocrine & Metabolic
Abilify - <i>ariprazole</i> Ativan - <i>lorazepam</i> Clozaril - <i>clozapine</i> Depakote - <i>divalproex</i> Haldol - <i>haloperidol</i> Invega - quetiapine Klonopin - <i>clonazepam</i> Lamictal - <i>lamotrigine</i> Lithium - <i>eskalith</i> Risperdal - Seroquel - Tripleptal Zyprexa -	Advair Discus Atrovent Flonase Flovent Loratadine Proventil	Crestor - <i>rosuvastatin</i> Lipitor Zocor	Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid
Seizures	Gastrointestinal Disorder Constipation	Blood Pressure Meds	Pain & Inflammation
Dilantin Keppra Neurontin Topamax	Colace - <i>docosate sodium</i> Miralax Prilosec Protonix Zantac 360 - <i>famotidine</i>	HCTZ (hydorchlorothiazide) Lisinopril Toprol Tenormin - <i>atenolol</i>	Flexeril Motrin Norco Tylenol with Codeine Ultram

9574
5238
1763

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 9574 DMA Code #2 5238 DMA Code #3 1763

DRUG NAME	Ativan
GENERIC NAME	lorazepam
DOSAGE RANGE	0.5- 4 mg Adult
HOW TO TAKE	by pill-whole: do not crush or chew. liquid form taken by mouth
USES	anxiety, insomnia, status epilepticus, nausea & vomiting - <u>chlorpromazine</u> (ref: dysreg)
SIDE EFFECTS	Sedation, dizziness, ataxia, respiratory depression, fatigue, appetite change
WARNINGS	misuse can cause addiction, can slow or stop breathing, can cause overdose & death

DRUG NAME	Ativan Klonopin
GENERIC NAME	Clonazepam
DOSAGE RANGE	0.5 mg, 1mg, 2mg
HOW TO TAKE	Swallow tablet with full glass of water
USES	Anxiety, panic disorder, seizure disorder, restless leg syndrome, REM sleep behavioral disorder
SIDE EFFECTS	Drowsiness, dizziness, impaired coordination, depression, fatigue, constipation
WARNINGS	Black box warning - addiction, abuse & misuse, dependence & withdrawal, appetite changes reactions

DRUG NAME	Abilify
GENERIC NAME	aripiprazole
DOSAGE RANGE	2mg, 5mg, 10mg, 15mg, 20mg, 30mg
HOW TO TAKE	whole, by mouth. Do not chew or crush tablet. do not spit out the oral disintegrating tab.
USES	Bipolar disorder, major depressive disorder, schizophrenia
SIDE EFFECTS	Blurred vision, increased salivation, uncontrolled muscle movements, headache, dizziness, drowsiness, weight changes
WARNINGS	Dementia related psychosis, suicidality

DRUG NAME	Clozaril
GENERIC NAME	Clozapine
DOSAGE RANGE	25mg, 100mg
HOW TO TAKE	oral liquid - use dosing syringe, orally disintegrating tablet - do not chew
USES	Schizophrenia, suicide prevention
SIDE EFFECTS	dyslipidemia, hyperglycemia, weight gain, drowsiness, constipation, dizziness, headache
WARNINGS	Severe neutropenia, orthostatic hypotension, bradycardia, seizures, dementia, psychosis

DRUG NAME	Crestor
GENERIC NAME	rosuvastatin
DOSAGE RANGE	5mg, 10mg, 20mg, 40mg
HOW TO TAKE	as directed, with or without food.
USES	hypercholesterolemia, mixed dyslipidemia, hypertriglyceridemia, dysbetalipoproteinemia <small>cardiovascular event prevent.</small>
SIDE EFFECTS	headache, myalgia, abdominal pain, nausea, dizziness, constipation, insomnia
WARNINGS	Do not take while pregnant, have liver problems, kidney disease, thyroid disorder

DRUG NAME	Apidra
GENERIC NAME	insulin glulisine
DOSAGE RANGE	U-100 pen, U-100 vial Uunits per mL
HOW TO TAKE	0.5-1 units Less than 15 minutes before or 20 min after meals.
USES	diabetes mellitus
SIDE EFFECTS	hypoglycemia, seizures, hypokalemia, hypersensitivity, anaphylaxis
WARNINGS	

DRUG NAME	Colace
GENERIC NAME	docusate sodium
DOSAGE RANGE	50-150 mg/day for 2-11 yrs 50-300 mg/day for 12 and older
HOW TO TAKE	by mouth Tablet or liquid
USES	constipation
SIDE EFFECTS	diarrhea, abdominal cramps, throat irritation, rash, electrolyte disorders
WARNINGS	

DRUG NAME	HCTZ
GENERIC NAME	hydrochlorothiazide
DOSAGE RANGE	Cap: 12.5mg Tab: 12.5mg, 25mg, 50mg
HOW TO TAKE	By mouth
USES	high blood pressure, peripheral edema, renal dosing
SIDE EFFECTS	weakness, feeling like you may pass out, severe spreading stomach pain, electrolyte imbalance
WARNINGS	

DRUG NAME	
GENERIC NAME	
DOSAGE RANGE	
HOW TO TAKE	
USES	
SIDE EFFECTS	
WARNINGS	

DRUG NAME	
GENERIC NAME	
DOSAGE RANGE	
HOW TO TAKE	
USES	
SIDE EFFECTS	
WARNINGS	