

Darice Darby

**DMA TRAINING**  
**LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING**

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [ 48 ]  
 See slide 65 in DMA Packette

<b>Mental Illness Anxiety Disorders</b>	<b>Inhalers Allergy / Asthma</b>	<b>Hyperlipidemia Statins</b>	<b>Diabetes Endocrine &amp; Metabolic</b>
Abilify Ativan Clozaril ✓ Depakote Haldol ✓ Invega ✓ Klonopin ✓ Lamictal ✓ Lithium Risperdal Seroquel Tripleptal ✓ Zyprexa	Advair Discus Atrovent Flonase Flovent Loratadine Proventil	Crestor Lipitor Zocor	Apidra Byetta Glucophage ✓ Glyburide Lantus Levemir Levothyroxine ✓ Novolog Synthroid
<b>Seizures</b>	<b>Gastrointestinal Disorder Constipation</b>	<b>Blood Pressure Meds</b>	<b>Pain &amp; Inflammation</b>
Dilantin Keppra Neurontin Topamax ✓	Colace Miralax Prilosec Protonix ✓ Zantac	HCTZ (hydrochlorothiazide) Lisinopril Toprol Tenormin	Flexeril Motrin Norco Tylenol with Codeine Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 0611

DMA Code #2 0925

DMA Code #3 0515

DRUG NAME	Topamax
GENERIC NAME	Topiramate
DOSAGE RANGE	25mg-200mg
HOW TO TAKE	PO qd/qwk - by mouth daily/weekly
USES	Seizure disorder, PTSD, migraine, parasomnias
SIDE EFFECTS	tiredness, drowsiness, dizziness, diarrhea, loss of appetite
WARNINGS	may make you dizzy or impair judgment, may make you more likely to stroke / drink lots of fluid when hot, if fever get medical help

DRUG NAME	Protonix
GENERIC NAME	pantoprazole
DOSAGE RANGE	20 mg - 40 mg
HOW TO TAKE	PO qd - by mouth daily
USES	acid reflux, heartburn, difficulty swallowing
SIDE EFFECTS	headache, diarrhea, low blood level, rash, joint pain
WARNINGS	may increase risk for bone fractures, passes into breast milk

DRUG NAME	Levothyroxine
GENERIC NAME	Synthroid, Levoxyl
DOSAGE RANGE	25 mcg (.025 mg) - 500 mcg (.5 mg)
HOW TO TAKE	PO qd - by mouth daily
USES	over active thyroid, enlarged thyroid gland, diabetes
SIDE EFFECTS	weight gain/loss, headache, vomiting, diarrhea, fever, sensitivity to heat
WARNINGS	Not for obesity / weight loss

DRUG NAME	Glucophage
GENERIC NAME	metformin
DOSAGE RANGE	500 mg - 1000 mg
HOW TO TAKE	PO bid - by mouth twice a day with meals
USES	diabetes, control high blood sugar
SIDE EFFECTS	upset stomach, vomiting, diarrhea, nausea, weakness
WARNINGS	lactic acidosis

DRUG NAME	Tripleptal
GENERIC NAME	Oxcarbazepine
DOSAGE RANGE	150 mg - <del>600mg</del> 1200 mg
HOW TO TAKE	PO
USES	Seizures
SIDE EFFECTS	Suicidal behavior + ideation, hyponatremia, dizziness, vomiting, headache
WARNINGS	avoid abrupt withdrawal

DRUG NAME	Lamictal
GENERIC NAME	Lamotrigine
DOSAGE RANGE	25mg - 200mg
HOW TO TAKE	PO - ODT / chewable qd - daily
USES	seizures, bipolar disorder
SIDE EFFECTS	rash, dizziness, vertigo, headache, nausea, back pain, fever
WARNINGS	Serious rash requiring hospitalization

DRUG NAME	Klonopin
GENERIC NAME	Clonazepam
DOSAGE RANGE	.5mg - 2mg
HOW TO TAKE	PO bid-tid (2 or 3 x day)
USES	Seizure, panic disorder, anxiety, sleep terrors, sleepwalking
SIDE EFFECTS	drowsiness, depression, fatigue, confusion, rash, constipation
WARNINGS	risk from concomitant opioid use, Addiction, Abuse, misuse, dependence

DRUG NAME	Invega
GENERIC NAME	Paliperidone
DOSAGE RANGE	1.5mg - 9mg
HOW TO TAKE	PO qam (every morning)
USES	Schizophrenia, schizoaffective disorder, renal dosing
SIDE EFFECTS	headache, anxiety, dizziness, weight gain, cough, fatigue, abdominal pain
WARNINGS	Not approved for dementia-related psychosis

DRUG NAME	Haldol
GENERIC NAME	haloperidol
DOSAGE RANGE	.5mg - 20mg
HOW TO TAKE	PO bid-tid (2 or 3 x day)
USES	Psychosis, tourette syndrome, acute agitation
SIDE EFFECTS	insomnia, anxiety, drowsiness, lethargy, weight change, seizures
WARNINGS	Not approved for dementia-related psychosis

DRUG NAME	Clozaril
GENERIC NAME	Clozapine
DOSAGE RANGE	25mg - 100mg
HOW TO TAKE	PO bid (twice a day)
USES	Schizophrenia, suicide prevention, renal dosing
SIDE EFFECTS	Stroke, seizures, fever, nausea, headache, weight gain, tremor
WARNINGS	Severe neutropenia, seizures, dementia-related psychosis