

DMA TRAINING

LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
 See slide 65 in DMA Packette

Mental Illness Anxiety Disorders	Inhalers Allergy / Asthma	Hyperlipidemia Statins	Diabetes Endocrine & Metabolic
Abilify Ativan Clozaril Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa	Advair Discus Atrovent Flonase Flovent Loratadine Proventil	Crestor Lipitor Zocor	Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid
Seizures	Gastrointestinal Disorder Constipation	Blood Pressure Meds	Pain & Inflammation
Dilantin Keppra Neurontin Topamax	Colace Miralax Prilosec Protonix Zantac	HCTZ (hydrochlorothiazide) Lisinopril Toprol Tenormin	Flexeril Motrin Norco Tylenol with Codeine Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 1657

DMA Code #2 1253

DMA Code #3 9532

DRUG NAME	Abilify	
GENERIC NAME	aripiprazole	
DOSAGE RANGE	2-30 mg	
HOW TO TAKE	Dissolving tablet, liquid medicine by mouth	
USES	agitation, distress, restlessness, mania like symptoms, ticure	5
SIDE EFFECTS	Schizophrenia, bipolar I, major depressive disorder	
WARNINGS	Not approved for dementia related psychosis suicide risk for old people under 24 and over 65	

DRUG NAME	Ativan	
GENERIC NAME	lorazepam	
DOSAGE RANGE	0.5-2 mg Tabs / 2, 4 mg per mL injections	
HOW TO TAKE	1 tablet by mouth or IV injection	
USES	short term insomnia, status epilepticus / chemo related nausea	
SIDE EFFECTS	Sedation, dizziness, ataxia, resp depression, hypoventilation	
WARNINGS	Concomitant Opioid Use, Abuse/Addiction, Dependence, Withdrawal	

DRUG NAME	Haldol	
GENERIC NAME	Haloperidol	
DOSAGE RANGE	Tab: 0.5-20 mg	
HOW TO TAKE	Tab or liquid by mouth	
USES	psychosis, tourette syndrome, acute agitation	
SIDE EFFECTS	tardive dyskinesia, ataxia, insomnia, anxiety, drowsiness	
WARNINGS	Not approved for dementia related psychosis	

DRUG NAME	Lithium lithium	
GENERIC NAME	Lithium lithium	
DOSAGE RANGE	Cap: 150-600 mg, Tab: 30 mg, ER Tab: 300-450 mg SOL: 8 mEq per 5 ml	
HOW TO TAKE	Swallow Tab by Tab Cap or liquid by mouth	
USES	bipolar I maintenance / bipolar I disorder, schizoaffective, renal dosing	
SIDE EFFECTS	tremor, polyuria, polydipsia, weight gain, diarrhea	
WARNINGS	Lithium Toxicity	

DRUG NAME	Risperdal	
GENERIC NAME	risperidone	
DOSAGE RANGE	Tab: 0.25-4 mg, SOL 1 mg per mL	
HOW TO TAKE	dissolving tablet or liquid medicine by mouth	
USES	schizophrenia, bipolar I, tourette syndrome, renal dosing, hepatic dosing	
SIDE EFFECTS	increased appetite, fatigue, insomnia, nausea, cough, constipation	
WARNINGS	Not approved for dementia related psychosis	