

|              |  |
|--------------|--|
| DRUG NAME    | Ambility   |
| GENERIC NAME | Aripiprazole   |
| DOSAGE RANGE | 2mg-30mg tab   |
| HOW TO TAKE  | by mouth   |
| USES         | <del>schizo</del> schizophrenia, bipolar disorder, major depressive dis. |
| SIDE EFFECTS | seizures, anaphylaxis, stroke, syncope, dystonia                         |
| WARNINGS     | not for dementia patients; can cause suicidal thoughts or actions.       |

|              |  |
|--------------|--|
| DRUG NAME    | <del>Dilantin</del> Dilantin                           |
| GENERIC NAME | phenytoin  |
| DOSAGE RANGE | capsules 30mg, 100mg Chewable 50mg, Susp 125mg per 5ml |
| HOW TO TAKE  | I.V. or by mouth                                       |
| USES         | seizure disorder, epilepsy                             |
| SIDE EFFECTS | nausea, vomiting, anaphylaxis, dizziness, tremors.     |
| WARNINGS     | Cardiovascular Risk with rapid infusion                |

|              |   |
|--------------|---|
| DRUG NAME    | Colace  |
| GENERIC NAME | docusate sodium                                     |
| DOSAGE RANGE | 50-300mg  |
| HOW TO TAKE  | oral  |
| USES         | laxative for softening stool                        |
| SIDE EFFECTS | diarrhea, cramps abdominal, throat irritation, rash |
| WARNINGS     | nausea, vomiting, appendicitis, fecal impaction.    |

|              |   |
|--------------|---|
| DRUG NAME    | Toprol  |
| GENERIC NAME | metoprolol succinate  |
| DOSAGE RANGE | 25mg-200mg tab  |
| HOW TO TAKE  | oral  |
| USES         | HTN, angina, heart failure, migraines                         |
| SIDE EFFECTS | CHF, hepatitis, gangrene, photosensitivity, rash, depression. |
| WARNINGS     | avoid abrupt cessation.                                       |

|              |  |
|--------------|--|
| DRUG NAME    | Miralax  |
| GENERIC NAME | polyethylene glycol 3350                         |
| DOSAGE RANGE | 1 caplet   |
| HOW TO TAKE  | dissolve into water, take orally.                |
| USES         | constipation relief                              |
| SIDE EFFECTS | nausea, cramping, diarrhea, laxative dependence. |
| WARNINGS     | renal impairment, GI obstruction                 |

|              |  |
|--------------|--|
| DRUG NAME    | Crestor  |
| GENERIC NAME | rosuvastatin   |
| DOSAGE RANGE | 5mg - 40 mg TAB  |
| HOW TO TAKE  | oral   |
| USES         | hypercholesterolemia, mixed dyslipidemia, hypertriglyceridemia           |
| SIDE EFFECTS | myopathy, tendon rupture, headache, nausea, myalgia, acute renal failure |
| WARNINGS     | caution if hypothyroidism  |

|              |   |
|--------------|---|
| DRUG NAME    | Haldol  |
| GENERIC NAME | haloperidol                                   |
| DOSAGE RANGE | .5mg - 20mg tab                               |
| HOW TO TAKE  | oral by mouth                                 |
| USES         | Tourette syndrome, acute agitation, psychosis |
| SIDE EFFECTS | Sudden death, dystonia, seizures, leukopenia  |
| WARNINGS     | Dementia Related psychosis                    |

|              |   |
|--------------|---|
| DRUG NAME    | Keppra  |
| GENERIC NAME | levetiracetam   |
| DOSAGE RANGE | 250mg tab - 1000mg tab. Sol 100mg per ml, various injection.            |
| HOW TO TAKE  | renal, oral, injection.   |
| USES         | partial seizures, juvenile myoclonic epilepsy                           |
| SIDE EFFECTS | depression, hostility, psychosis, headache, pain, irritability, nausea. |
| WARNINGS     | NA  |

|              |  |
|--------------|--|
| DRUG NAME    | Zantac   |
| GENERIC NAME | <del>z</del> famotidine                                      |
| DOSAGE RANGE | 1-2 tab  |
| HOW TO TAKE  | oral, renal  |
| USES         | preventing heartburn   |
| SIDE EFFECTS | AV block, hepatitis, pneumonia, constipation, taste changes. |
| WARNINGS     | na   |

|              |  |
|--------------|--|
| DRUG NAME    | Byetta   |
| GENERIC NAME | exenatide  |
| DOSAGE RANGE | 5mcg - 10mcg inj   |
| HOW TO TAKE  | injection per, oral, renal   |
| USES         | diabetes mellitus, type 2  |
| SIDE EFFECTS | nausea, vomiting, <sup>hypersensitivity</sup> sensitivity, acute kidney injury, headache |
| WARNINGS     | na.  |