

Tiara Pollard
Nov. 1, 2021

DMA TRAINING
LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
See slide 65 in DMA Packette

Mental Illness Anxiety Disorders	Inhalers Allergy / Asthma	Hyperlipidemia Statins	Diabetes Endocrine & Metabolic
Abilify Ativan Clozaril Depakote Haldol Invega <u>Klonopin</u> Lamictal Lithium <u>Risperdal</u> Seroquel Tripleptal Zyprexa	Advair Discus Atrovent <u>Flonase</u> <u>Flovent</u> Loratadine Proventil	<u>Crestor</u> <u>Lipitor</u> Zocor	Apidra <u>Byetta</u> <u>Glucophage</u> Glyburide <u>Lantus</u> Levemir <u>Levothyroxine</u> Novolog Synthroid
Seizures	Gastrointestinal Disorder Constipation	Blood Pressure Meds	Pain & Inflammation
Dilantin <u>Keppra</u> <u>Neurontin</u> Topamax	Colace <u>Miralax</u> Prilosec Protonix <u>Zantac</u>	HCTZ (hydrochlorothiazide) <u>Lisinopril</u> Toprol Tenormin	Flexeril Motrin Norco <u>Tylenol with Codeine</u> Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 0611

DMA Code #2 0925

DMA Code #3 0615

DRUG NAME	Byetta
GENERIC NAME	Exenatide
DOSAGE RANGE	250mcg (5mg per dose, 60 doses) (10mg per dose, 60 doses)
HOW TO TAKE	inject this medication under skin in the thigh, abdomen, or upper arm.
USES	help control blood sugar levels in adults with type 2 diabetes
SIDE EFFECTS	vomiting, diarrhea, nervousness, or upset stomach.
WARNINGS	you should not use this if you have severe kidney disease

DRUG NAME	Miralax
GENERIC NAME	Polyethylene glycol
DOSAGE RANGE	17g (powder)
HOW TO TAKE	Stir and dissolve one packet of powder in any 4 to 8 ounce drink
USES	Used to treat occasional constipation
SIDE EFFECTS	rectal irritation, malaise, nausea, vomiting, dehydration, polydipsia
WARNINGS	Stop taking if you have severe or bloody diarrhea; rectal bleeding

DRUG NAME	Keppra
GENERIC NAME	Levetiracetam
DOSAGE RANGE	250mg (blue) 500mg (yellow) 750mg (orange) 1000mg (white)
HOW TO TAKE	by mouth once daily
USES	Used to treat the symptoms of partial onset seizures, tonic-clonic seizures
SIDE EFFECTS	Drowsiness, Dizziness, Unusual tiredness, or weakness may occur.
WARNINGS	You should not stop taking Keppra without consulting your doctor

DRUG NAME	Tylenol with Codeine
GENERIC NAME	Acetaminophen-codeine
DOSAGE RANGE	15 to 60 mg 360 mg in 24 hours
HOW TO TAKE	Orally up to every 4 hours
USES	help to relieve mild to moderate pain. it contains an opioid-non-opioid ^{pain} reliever.
SIDE EFFECTS	hyperhidrosis, sedation, constipation, drowsy, vomiting, nausea
WARNINGS	May cause drowsiness and dizziness. Alcohol and marijuana may intensify.

DRUG NAME	Lisinopril
GENERIC NAME	Prinivil
DOSAGE RANGE	10mg 20mg
HOW TO TAKE	by mouth
USES	used to treat high blood pressure in adults and children 4 years or older.
SIDE EFFECTS	dizziness, headache, URI, fatigue, cough...
WARNINGS	This drug may impair the ability to operate a vehicle, vessel, or machinery

DRUG NAME	Klonopin
GENERIC NAME	Clonazepam
DOSAGE RANGE	0.5mg, 1mg, 2mg
HOW TO TAKE	By mouth
USES	Anxiety, Confusional arousals, sleep walking
SIDE EFFECTS	Muscle weakness, Drowsy
WARNINGS	May cause drowsiness + dizziness. Alcohol-marijuana may intensify this effect. Use car when operating a vehicle, vessel, or machinery

DRUG NAME	Risperdal
GENERIC NAME	Risperidone
DOSAGE RANGE	0.25mg, 0.5mg, 1mg, 2mg, 3mg, 4mg
HOW TO TAKE	By mouth (with or without food)
USES	Schizophrenia, Bipolar
SIDE EFFECTS	extrapyramidal effects, dizziness, dry mouth, fever, weight gain.
WARNINGS	Not approved in elderly patients and can cause death

DRUG NAME	Fluvent
GENERIC NAME	Fluticasone
DOSAGE RANGE	44mcg, 110mcg, 220mcg, 88mcg
HOW TO TAKE	Shake 5 to 10 sec. push to down one time while breathing in hold for 5-10 sec. breathe out.
USES	Control and prevent symptoms caused by asthma.
SIDE EFFECTS	Hoarseness or throat irritation may occur.
WARNINGS	fungal infections in you mouth or throat. Weaken immune system

DRUG NAME	Lipitor
GENERIC NAME	Atorvastatin
DOSAGE RANGE	10mg, 20mg, 40mg, 80mg
HOW TO TAKE	by mouth once daily
USES	is used along with a proper diet to help lower 'bad'
SIDE EFFECTS	Constipation, diarrhea, nausea, fatigue, gas, heart burn, headache, muscle pain
WARNINGS	Do not use if you are allergic to, or if you have liver disease.

DRUG NAME	Lantus
GENERIC NAME	is only available in brand version
DOSAGE RANGE	100 units, 300 units
HOW TO TAKE	take by mouth once a day.
USES	is a long acting insulin that helps control high blood sugar levels.
SIDE EFFECTS	Weight gain, Allergic reaction, Diarrhea, Common Cold, Swelling, Inflammation...
WARNINGS	Never share lantus pens, syringes, or needles with any one else.