



Medication Administration In-Service and Evaluation

Name of Facility/Home: Trolley house - coopersville.

Employee Receiving In-Service: Kassandra J Wydeck

Date of 1st In-Service: 10 / 14 / 2021 Time: 1 : 00pm am / pm Trainer: Learning and Development

Date of 2nd In-Service: 10 / 14 / 2021 Time: 3 : 00pm am / pm Trainer: Learning and Development

Date of 3rd In-Service: 10 / 21 / 2021 Time: 7 : 30 am / ~~pm~~ Trainer: Kory Bickford

Date of 4th In-Service: 10 / 22 / 21 Time: 6 : 20 am / pm Trainer: Kory Bickford

Date of 5th In-Service: 10 / 24 / 21 Time: 7 : 30 am / pm Trainer: Harry Snyder

Date of 6th In-Service: 10 / 26 / 21 Time: 6 : 30 ^{Am} / pm Trainer: Harry Snyder

Date of Final Evaluation: 10 / 27 / 21 Time: 6 : 45 am / pm Trainer: [Signature]

All staff must complete all three (6) In-Services and Final Evaluation

Instructions: Check (✓) the appropriate box after Employee has been in-serviced.

| | | In-Service # | 1st | 2nd | 3rd | 4th | 5th | 6th | Eval. | Comments |
|---|--|--------------|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|----------|
| 1 | Medication Area | | <input checked="" type="checkbox"/> | |
| | a. Location of ample supplies prior to administration | | <input checked="" type="checkbox"/> | |
| | b. Area is clean and organized | | <input checked="" type="checkbox"/> | |
| | c. Area is always locked | | <input checked="" type="checkbox"/> | |
| | d. Location of all medication: Internal, External, Refrigerated, Controlled Drugs (narcotics) | | <input checked="" type="checkbox"/> | |
| 2 | DMA washes hands prior to administering medications and between each Resident | | <input checked="" type="checkbox"/> | |
| 3 | Medication keys are retained by DMA | | <input checked="" type="checkbox"/> | |
| 4 | Resident is identified per facility policy and procedure prior | | <input checked="" type="checkbox"/> | |
| 5 | Vital signs are taken per facility policy prior to administering medications (if applicable), always on cardiac and BP medications | | <input checked="" type="checkbox"/> | |
| | a. If Pulse and BP are required, hands and equipment are washed per facility policy | | <input checked="" type="checkbox"/> | |
| | b. If Apical Pulse is required, privacy is provided | | <input checked="" type="checkbox"/> | |
| 6 | Medications Administration per facility policy and procedure: to include review of the '6 Rights' | | <input checked="" type="checkbox"/> | |
| | a. Medications are properly removed from container/blister pack and (.) dot is placed in appropriate box on MAR | | <input checked="" type="checkbox"/> | |
| | b. Liquid medication is poured at eye level, with palm covering label of stock bottle | | <input checked="" type="checkbox"/> | |



Medication Administration In-Service and Evaluation

| In-Service # | | 1st | 2nd | 3rd | 4th | 5th | 6th | Eval. | Comments |
|---|---|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|-------------------------------------|----------|
| 6 | c. DMA verifies medication and strength with order as transcribed on medication record per facility policy and procedure | <input checked="" type="checkbox"/> | |
| | d. Observe Resident to ensure medication is swallowed | <input checked="" type="checkbox"/> | |
| | e. Offer adequate and appropriate fluid with medication | <input checked="" type="checkbox"/> | |
| | f. Medication record is signed immediately after administration of same | <input checked="" type="checkbox"/> | |
| | g. Controlled substance record is signed immediately after administration of same | <input checked="" type="checkbox"/> | |
| | h. Correct dose is administered | <input checked="" type="checkbox"/> | |
| | i. Medication is administered at correct time | <input checked="" type="checkbox"/> | |
| j. Verify no additional MAR pages have been added | <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/> | | |
| 7 | Infection control technique is reviewed | <input checked="" type="checkbox"/> | |
| 8 | Medication via gastric tube administered per facility policy and procedure (if applicable) | <input checked="" type="checkbox"/> | |
| | a. Resident is properly positioned, at a 45° sitting angle | <input checked="" type="checkbox"/> | |
| | b. Tube is checked for placement and patency | <input checked="" type="checkbox"/> | |
| | c. Tube is flushed before, between and after medications are administered | <input checked="" type="checkbox"/> | |
| 9 | Injections are administered by the Resident or a DMA if there is a doctor's order present, per facility policy and procedure | <input checked="" type="checkbox"/> | |
| | a. Syringes and needles are disposed of in sharps container, by person giving the injection without recapping | <input checked="" type="checkbox"/> | |
| | b. Proper glucometer testing is observed. Determination of competence re: accurately perform and read glucometer testing results | <input checked="" type="checkbox"/> | |
| 10 | DMA crushes medication according to facility policy and procedure ONLY with physician's orders. | <input checked="" type="checkbox"/> | |
| 11 | DMA administers eye and ear medication according to facility policies and procedures | <input checked="" type="checkbox"/> | |
| 12 | Side effects of psychoactive medication are noted (lethargy, hallucinations) and reported. | <input checked="" type="checkbox"/> | |
| 13 | Medication administration should not interrupted. DO NOT RUSH | <input checked="" type="checkbox"/> | |
| 14 | Controlled drugs are stored (Double Locked) according to facility policy and procedure | <input checked="" type="checkbox"/> | |
| 15 | Residents' rights are observed | <input checked="" type="checkbox"/> | |
| 16 | Location, Procedures and Documenting for administering PRN | <input checked="" type="checkbox"/> | |
| 17 | Designated Medication Administrator follows facility policy and procedure for medications refused or withheld. (MER & IR written) | <input checked="" type="checkbox"/> | |
| 18 | Medications are administered within time frame per facility policy | <input checked="" type="checkbox"/> | |



Medication Administration In-Service and Evaluation

| | In-Service # | 1st | 2nd | 3rd | 4th | 5th | 6th | Eval. | Comments |
|----|---|-----|-----|-----|-----|-----|-----|-------|----------|
| 19 | Medication errors are reported to Site Supervisor and RN teaching medication classes | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 20 | Medication area is cleaned and locked after completion of medication administration | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 21 | Designated Medication Administrator can identify action and common side effects of medications administered | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 22 | Approved Abbreviations List is reviewed | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 23 | Seizure precautions and documentation | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 24 | After hour procedures, procedures for found/spilled medication, location of Guide to Drugs Book | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 25 | 2nd Staff Verification, what it is, when it is needed, and how to document it | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |
| 26 | Refusal of Medication procedures (prompt 3 times, then write appropriate documentation) | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | ✓ | |

FOLLOW UP CONCERNS

Specify time frame for completion: _____ N/A

I have received the above In-service and have read the Organizations **Medical Policies**. I understand what is expected of me as a Designated Medication Administrator. I also understand that any immediate medical questions or concerns should be directed to the Coordinator of Care at my Site during open office hours and to the On-Call person after hours.

Rassanara Wydeck
Employee Signature

10-24-21
Date

[Signature]
Home Manager Signature

10-24-21
Date

DMA TRAINING

LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
See slide 65 in DMA Packette

| Mental Illness Anxiety Disorders | Inhalers Allergy / Asthma | Hyperlipidemia Statins | Diabetes Endocrine & Metabolic |
|--|--|--|---|
| Abilify Ativan Clozaril Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa | Advair Discus Atrovent Flonase Flovent Loratadine Proventil | Crestor Lipitor Zocor | Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid |
| | | | |
| Seizures | Gastrointestinal Disorder Constipation | Blood Pressure Meds | Pain & Inflammation |
| Dilantin Keppra Neurontin Topamax | Colace Miralax Prilosec Protonix Zantac | HCTZ (hydrochlorothiazide) Lisinopril Toprol Tenormin | Flexeril Motrin Norco Tylenol with Codeine Ultram |

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 9532

DMA Code #2 1737

DMA Code #3 1978

10 drug name: Risperdal
generic name: risperidone
dosage: TAB: 0.5mg 1mg 2mg 3mg 4mg SOL 1mg per mL
How to take: orally / by mouth 1-2 times a day
uses: Schizophrenia, bipolar I disorder,
Tourette Syndrome

Side effects: severe hypotension, stroke, hyperglycemia, appetite increase, cough, vomiting, fever, anxiety, dizziness, rash, confusion, abdominal pain,
Warnings: not approved for dementia-related psychosis. most deaths due to cardiovascular or infectious events.

11 drug name: Seroquel
generic name: quetiapine
dosage: TAB: 25mg 50mg 100mg 200mg 300mg 400mg
how to take: by mouth - daily - 3x

uses: Schizophrenia, bipolar I disorder, bipolar disorder, generalized anxiety disorder
Side effects: cataracts, pancreatitis, suicidality, anaphylaxis, dizziness, rash, anemia, headache, fatigue, fever, insomnia, nausea, fever, vomiting.
Warnings: Dementia-related psychosis, not approved. Suicidality risk in children, adolescents, and young adults w/ major depressive or other psychiatric disorders.

drug name: Loratadine

generic name: Loratadine

dosage: TAB: 10mg ODT: 10mg SOL: 1mg per mL

How to take: by mouth / daily.

uses: allergy, chronic idiopathic urticaria,

side effects: bronchospasm, anaphylaxis, seizures, drowsiness, fatigue, abdominal pain, headache, diarrhea.

warnings: do not take this medicine if you are allergic to loratadine. Can pass into breast milk and may harm a nursing baby. Tell dr. if you're breastfeeding, if you have allergies or medical conditions.

drug name: Atrovent HFA

generic name: ipratropium bromide inhaled

dosage: MDI 17mcg per actuation

how to take: Puff every 20 min as needed up to 3h

uses: COPD, maintenance, asthma exacerbation,

side effects: hypersensitivity, anaphylaxis, bronchospasm, paradoxical, glaucoma, nausea, sinusitis, dizziness, back pain, urinary retention, bronchitis, UTI

warnings: do not get in eyes, not approved for anyone under 18 yr (aerosol) under 12 yr (solution)

18
Drug name: Flovent HFA
Generic name: fluticasone propionate inhaled
Dosage: MDI - 44 mcg per actuation, 110 mcg, 220 mcg per act.
How to: puffs inhaled daily.
Uses: Asthma.

Side effects: glaucoma long term, cataracts long-term, anaphylaxis, headache, throat irritation, cough, nausea, vomiting, rash, hearseness.
Warnings: do not use to treat asthma attack that has already begun. long term use may lead to bone loss. tellor if you are pregnant.

19
Drug name: Flovent Diskus
Generic name: fluticasone propionate inhaled
Dosage: DPI: 100 mcg per blister, 250 mcg per blister
How to: puffs daily.
Uses: Asthma maintenance.

Side effects: anaphylaxis, cataracts (long term), growth suppression, bronchospasms, headache, throat irritation, cough, nausea, rash, hoarseness.
Warnings: may not be safe to breastfeed, do not use if allergic, do not use to treat an asthma attack that has already begun. do not use inhalation powder if you are allergic to milk proteins.

22
drug name: Lipitor
generic name: atorvastatin.
dosage: TAB: 10mg 20mg 40mg 80mg.
how to take: by mouth daily.
uses: hypercholesterolemia. mixed dyslipidemia.
hypertriglyceridemia, type IV, dysbetalipoproteinemia.
homozygous familial hypercholesterolemia.
cardiovascular event prevention. Atherosclerotic
cardiovascular dz.
side effects: myopathy. acute renal failure.
tendon rupture. UTI. diarrhea. extremity pain.
muscle spasm. insomnia. cognitive impairment.
warnings: do not take if pregnant or
breastfeeding. Can cause muscle tissue breakdown
which can lead to kidney failure. Tell dr. about
all medications including ones you've just
started or stopped.

23
drug name: Zocor
generic name: Simvastatin
dosage: TAB 5mg 10mg 20mg 40mg 80mg
how to take: daily every evening.
uses: hypercholesterolemia. mixed dyslipidemia.
hypertriglyceridemia type IV. dysbetalipoproteinemia.
homozygous familial hypercholesterolemia.
cardiovascular event prevention. Atherosclerotic
cardiovascular dz

benztropine

generic.

26

TAB: 0.5mg 1mg 2mg INJ: 1mg per mL
by mouth at bedtime or INJ.
Parkinsonism, extrapyramidal sx,
dystonic rxns, acute.

tachycardia, psychosis, heart stroke
constipation, sedation, anorexia, rash, dizziness,
blurred vision.

do not use on persons under 3.
Older adults may be more sensitive. Tell your
dr if you are or may become pregnant.

Lorazepam

generic.

27

TAB: 0.5mg 1mg 2mg SOL 2mg per mL INJ: 2mg per mL
4mg per mL
daily 2-3x as needed for anx.
by mouth daily at bedtime as needed for insomnia.

Intravenous for status epilepticus. daily for nausea
intramuscular for pre-op sedation. PO/IM/IV every
hour as needed for alcohol withdrawal. IV every
5-15 mins as needed for delirium tremens.
by mouth for neuroleptic malignant syndrom.

respiratory depression, apnea,
dependency, abuse, jaundice, sedation, amnesia,
incontinence, constipation, menstrual irregularities.

may lead to dependence and/or abuse.
benzo use. may lead to sedation, res depression

drug name: Abilify

generic name: aripiprazole

dosage range: TAB: 2mg, 5mg, 10mg, 15mg, 20mg, 30mg

how to take: orally / by mouth - daily.

uses: schizophrenia, bipolar I disorder, manic depression.

side effects: stroke, pancreatitis, anaphylaxis, seizures, headache, nausea, weight gain, cough, insomnia, compulsive behaviors, fatigue, vomiting, anxiety, tremor

warnings: Dementia-related psychosis - not approved.

Increased mortality in elderly on conventional or atypical antipsychotics, most deaths due to cardiovascular or infectious events. not approved for ped's patients, increased suicidality in children, young adults w/ major depressive or other psychotic disorders.

drug name: Ativan

generic name: lorazepam

dosage range: TAB: .5mg, 1mg, 2mg (IV): 2mg per mL, 4mg per mL

how to take: orally, intravenous, intramuscular

uses: anxiety, insomnia, alcohol withdrawal, pre op sedation, chemo-related nausea/vomiting.

side effects: apnea, abuse, seizures, jaundice, suicidality, sedation, amnesia, irritability, libido changes, incontinence, menstrual irregularities

warnings: addiction, misuse, abuse, dependence & withdrawal reactions, concomitant opioid use may lead to respiratory depression and death.

#5 drug name: Haldol
generic name: Haloperidol
discontinued in U.S.

#6 drug name: Invega dosage: ER TAB: 1.5mg, 3mg, 6mg, 9mg
generic name: Paliperidone
how to take: by mouth every morning.

uses: schizophrenia, schizoaffective disorder

side effects: arrhythmia, hyperthermia, neutropenia, seizures, headache, anxiety, constipation, dizziness, abdominal pain, sleep disorder

warnings: not approved for dementia related psychosis, inc. mortality risk in elderly dementia pts on conventional or atypical antipsychotics. most deaths due to cardiovascular or infectious events.

#7 drug name: Klonopin
generic name: clonazepam

dosage: TAB 0.5mg, 1mg, 2mg

how to take: by mouth, 3x daily.

uses: panic disorder, anxiety, restless leg syndrome, confusion, arousal, sleep terrors, sleepwalking, rapid eye movement sleep behavior disorder, burning mouth syndrome, tourette syndrome

side effects: respiratory depression, abuse, seizures, dependency, drowsiness, depression, fatigue, amnesia, confusion, constipation, libido changes, rash.

warnings: concomitant opioid use (profound sedation, coma, death. assess risk for abuse, misuse, and addiction before prescribing, dependence and withdrawal
associated with significant physical dependence.

#8

drug name: Lamictal

generic name: Lamotrigine

dosage: TAB: 25mg, 100mg, 150mg, 200mg, CHEW: 2mg, 5mg, 25mg

how to take: orally / by mouth

uses: bipolar I disorder, partial seizures, seizures-Lennox-Gastaut Syndrome, seizures-Primary Generalized tonic clonic, migraine w/ aura prophylaxis

side effects: severe rash, suicidality, angioedema, dizziness, nausea, vertigo, rash, back pain, blurred vision, fever, vaginitis, anorexia, anxiety, irritability, speech disturbance, constipation, hot flashes, neck pain,

warnings: Serious rash requiring hospitalization, Stevens-Johnson Syndrome, rare cases of toxic epidermal necrolysis, rash related deaths.

#9

drug name: Lithium

generic name: Lithium

dosage: CAP: 150mg, 300mg, 600mg, TAB: 300mg, ER: 300mg SOL: 450mg, 8meq per 5ml

how to take: orally / by mouth 2-3 times a day

uses: bipolar I disorder maintenance, bipolar I disorder acute manic/mixed, schizoaffective disorder

side effects: coma, seizures, ventricular arrhythmic, severe bradycardia, weight gain, vomiting, drowsiness, anorexia, blurred vision, rash, acne, edema.

warnings: Lithium toxicity - close to therapeutic levels.

#3

drug name: Clozani

generic name: Clozapine

dosage range: TAB: 25mg; 100mg

how to take: by mouth, 2 times a day

uses: Schizophrenia; Suicide prevention; Schiz^{assoc}

side effects: Severe constipation, heat stroke, stroke, pulmonary embolism, myocarditis,

drowsiness, tremor; Constipation, fever, vomiting, confusion

warnings: must enroll in program, can

cause severe neutropenia, rapid dose escalation

can cause orthostatic hypotension; bradycardia

and syncope; seizures with first dose,

myocarditis, cardiomyopathy, mitral valve

incompetence, not approved for dementia

related psychosis

#4 drug name: Depakote

generic name: divalproex sodium

dosage: TAB: 125mg, 250mg, 500mg

how to take: by mouth / 2-3 times a day

uses: migraine headaches, partial seizures, absence seizures, bipolar disorder

side effects: bleeding, coma, hallucinations, headache, nausea, tremor, amnesia, depression, constipation, vision changes, nervousness

warnings: Hepatotoxicity, fetal risk (major)

congenital malformations, migraines in pregnant women, do not use for epilepsy or bipolar in pregnant or planning to become

28

drug name: cephalexin (Keflex)

generic name: generic.

dosage: CAP: 250mg 500mg 750mg TAB: 250mg 500mg
SUSP: 125mg per 5ml, 250mg per 5ml

uses: Infections, pharyngitis, UTI,
Skin/skin structure infections, endocarditis
Prophylaxis, dental.

route: by mouth. every 6-12 hours
for infections, UTI, every 12 hrs for Pharyngitis

side effects: hypersensitivity. anaphylaxis.
Cholestatic jaundice. diarrhea. nausea.
vomiting. abdominal pain.

warnings: may contain sugar. tell dr.
if you are pregnant or breast feeding.

29

drug name: ofloxacin.

generic name: generic.

dosage: TAB. 200mg 300mg 400mg.

route: infections (po q12h) bacterial
infections chlamydial (po q12h x 7 days)
urethritis, non-gon (po q12h x 7 days)
Prostatitis (po q12h x 6wk) UTI (po q12h x 10 days) or 3-7 days
epididymitis (po q12h x 10 days) leprosy (po qd
multibacillary) x 12mo.
leprosy paucibacillary (po qd x 24m) typhoid fever (po bid 7-14 days)

warnings & side effects: dizziness, nausea, anxiety, rash, vomiting, anaphylaxis, suicidality, depression, aortic dissection.

disabling potentially irreversible serious
reactions: avoid in myasthenia gravis.

#23
cont.

Side effects: myopathy, tendon rupture, acute renal failure, anaphylaxis, constipation, headache, eczema, vertigo, insomnia, gastritis.
Warnings: do not take while breastfeeding, or while pregnant. Stop right away and call dr. if you become pregnant. Can cause breakdown of skeletal ^{muscle} tissue leading to kidney failure.

#24

drug name: Apidra.
generic name: Insulin glulisine.
dosage: (INJ): (solostar U-100 pen) 100 units per mL (U-100 vial) 100 units per mL
how to take: two - four times daily.
uses: diabetes mellitus
Side effects: hypoglycemia, anaphylaxis, seizures, rash, weight gain, headache, influenza.

#25
~~26~~

drug name: ~~Colace~~ Colace
generic name: docusate sodium
dosage: 50-300mg/day - 12 and up.
how to take: by mouth daily/two times per day
uses: emollient laxative for stool softening
Side effects: diarrhea, abdominal cramps, throat irritation, rash, electrolyte disorders, bitter taste.

#20
drug name: Proventil HFA
generic name: albuterol inhaled
dosage: MDI: 90mcg per actuation
how to take: puff inhaled every 4-6 hrs prn
uses: bronchospasm, acute, exercise-induced
side effects: cardiac arrest, bronchospasm,
hypotension, bad taste, cough, pain, vomiting,
nervousness, tremor, tachycardia
warnings: do not use if allergic to milk
proteins, tell dr if you are pregnant,

#21
drug name: Crestor
generic name: rosuvastatin
dosage: TAB: 5mg 10mg 20mg 40mg
how to take: by mouth daily.
uses: hypercholesterolemia. mixed dyslipidemia
hypertriglyceridemia, type IV. dysbetalipoproteinemic
homozygous familial hypercholesterolemia
cardiovascular event prevention. atherosclerotic
cardiovascular dz
side effects: myopathy. tendon rupture.
acute renal failure. pancreatitis. headache.
abdominal pain. nausea. constipation. insomnia.
cognitive impairment.
warnings: Do not take while pregnant
or breast-feeding, if you have liver disease.
Can cause the breakdown of muscle tissue
which can lead to kidney failure. people of
asian descent may absorb at a higher
rate so discuss with dr to start a lower
starting dose.

#16

drug name: Atrovent Nasal
generic name: ipratropium bromide nasal
dosage: Spray (0.03%) 21 mcg per actuation (0.06%)
42 mcg per actuation
how to take: nostril 2x-3x a day.
uses: rhinorrhea (perennial allergic or nonallergic rhinitis, seasonal allergic rhinitis, common cold).
side effects: hypersensitivity, anaphylaxis, headache, nasal dryness, dry mouth, dry throat, taste changes, nasal irritation, nasal sx, nausea.
warnings: may pass into breastmilk, should not be used by child younger than 5 yr old

#17

drug name: Flonase Allergy relief
generic name: fluticasone propionate nasal
dosage: 1-11y - 1 actuation in each nostril daily - max 1
12y or older - 2 actuations each nostril daily.
how to: puff in each nostril (adults max 2, 2 each max 4)
uses: nasal polyp. allergy sx. Corticosteroid for relieving allergic rhinitis symptoms
side effects: cataracts (long-term) glaucoma, adrenal suppression, headache, nasal burning, nausea, cough, vomiting, abdominal pain, bronchitis, nasal ulcer,
warnings: use cautions.

#12

drug name: Zyprexa.

generic name: Olanzapine.

dosage: TAB: 2.5mg, 5mg, 7.5mg, 10mg, 15mg, 20mg ^{1NJ} 5mg per

How to take: by mouth, daily.

Uses: Schizophrenia, bipolar I disorder, agitation-
Schizophrenia or bipolar I disorder, major depressive
disorder, chemo-related nausea/vomiting,
refractory nausea/vomiting chemo-related

Side effects: Stroke, seizures, pancreatitis,
aspiration, insomnia, extremity pain, edema, tremor
constipation, appetite increase.

warnings:

not approved for dementia related psychosis.

#13

drug name: Advair Diskus

generic name: fluticasone propionate/salmeterol inhaled

dosage: DPI-Inhaler 100mcg/50mcg per blister,

250 mcg/50 mcg blister, 500 mcg/50mcg

How to take: 1 puff every 12 hrs.

Uses: Asthma, COPD

Side effects: pneumonia, bronchospasm, cardiac
arrest, immunosuppression (long term) headache,
sinusitis, nausea, vomiting, bronchitis, throat
irritation, dermatitis

warnings: Not a rescue medication, will not
work fast enough for asthma or bronchospasm
attack. seek med attention if you have
worsening breathing problems.