

GENERIC NAME	Clonazepam
DOSAGE RANGE	Tablet 0.25 mg
HOW TO TAKE	Oral Take two or three times as directed by doctor
USES	Use to prevent/control seizures
SIDE EFFECTS	Drowsiness, dizziness, tremors, loss of coordination.
WARNINGS	Caused dizziness/drowsiness.

DRUG NAME	Lamictal
GENERIC NAME	Lamotrigine
DOSAGE RANGE	Tablet 25mg, 100mg, 150mg, 200mg
HOW TO TAKE	By mouth as directed by doctor
USES	for bipolar disorder (manic depression)
SIDE EFFECTS	hives, blistering, peeling sores in your mouth and
WARNINGS	life-threatening skin rash, headache, vomiting

DRUG NAME	Lithium
GENERIC NAME	Lithium Carbonate
DOSAGE RANGE	300mg 450mg capsule
HOW TO TAKE	Taken by mouth
USES	by mouth
SIDE EFFECTS	Muscle weakness weight gain, & other side effects
WARNINGS	Vomiting severe hand tremors, vision change unsteadiness while standing or walking

DRUG NAME	Risperdal
GENERIC NAME	Risperidone
DOSAGE RANGE	2-3 mg once a day
HOW TO TAKE	Taken by mouth with or without food
USES	treat certain mental/mood disorders bipolar disorder
SIDE EFFECTS	dizziness, lightheadness, drooling, nausea, weight gain
WARNINGS	Increased or death elderly dementia patients.

DRUG NAME	Serpquel
GENERIC NAME	Quetiapine
DOSAGE RANGE	200 mg / 25 mg / 50 mg / 100 mg / 300 mg
HOW TO TAKE	Full glass of water With or without food
USES	By mouth 2 or 3 times a day.
SIDE EFFECTS	Constipation, drowsiness, upset stomach, tired
WARNINGS	

1422, 2725, 2066

GENERIC NAME	Ativan
DOSAGE RANGE	Lorazepam
HOW TO TAKE	One capsule once a day in the morning. Dose can be taken three times a day.
USES	
SIDE EFFECTS	Drowsiness, dizziness, loss of coordination, headache
WARNINGS	Nausea, blurred vision.

DRUG NAME	Clonazepam
GENERIC NAME	Latuda
DOSAGE RANGE	Treatment must begin at 12.5 mg
HOW TO TAKE	Take by mouth with or without food as directed
USES	Dose 12.5 mg once a day or twice daily.
SIDE EFFECTS	Weight gain, dizziness, fast heart rate, nasal dry mouth, fever, increased sweating.
WARNINGS	

DRUG NAME	Haldol
GENERIC NAME	Haloperidol decanoate
DOSAGE RANGE	10-20 times daily oral dose administered monthly
HOW TO TAKE	Two or three times a day.
USES	Get emergency
SIDE EFFECTS	Treatment of dementia, ICU agitation, psychosis, nausea
WARNINGS	Get emergency help if you have signs of an allergic

Reaction

DRUG NAME	Invega
GENERIC NAME	Paliperidone
DOSAGE RANGE	4 mg, 12 mg per day
HOW TO TAKE	by mouth with or without food
USES	Treat certain mental/mood disorders (such as schizophrenia)
SIDE EFFECTS	abnormal muscle, fast
WARNINGS	

DRUG NAME	Depakote
GENERIC NAME	Divalproex sodium
DOSAGE RANGE	
HOW TO TAKE	Take by mouth as directed by doctor.
USES	treat seizure disorders, control psychiatric conditions
SIDE EFFECTS	Diarrhea, dizziness, drowsiness, hair loss
WARNINGS	Warnings: Swollen feet, muscle, cramps