

How to Register for an Epocrates Account

Objective: Register for Epocrates.com

User Security Access Requirements: Available to all staff


Beacon User Role: Available to all staff

Timeline: Should create an account upon initiating DMA certification training process.

How to Register:

Pull up Google Chrome or any other web browser and go to <https://www.epocrates.com/>

You will now be the Epocrates home page.



Click on "Login" in the upper right-hand corner of the page.





A login box/registration box will pop-up.

The screenshot shows the Epocrates login and registration page. At the top, the Epocrates logo is displayed with the tagline "an athenahealth service". Below the logo, there is a text input field containing the email address "khaas@beaconspecialized.org". Underneath is a password field with masked characters. A checkbox labeled "Remember Me" is checked. A large black button with the text "Login" is positioned below the password field. Below the "Login" button, there are two links: "Forgot your login or password?" and "New User?". At the bottom of the form, there is another large black button with the text "Register Now".

Click on "Register Now" to register for an account. *(Enter credential if you have already registered to access the program.)*

Fill out the form below and click "Join". Use "Other Health Professional" for Occupation and use "49009" for your work zip. Be sure to use your Beacon e-mail.

The screenshot shows the "Create an Account" registration form. The form includes the following fields and options:

- Name:** First name and last name input fields.
- Work Email:** Input field for the user's work email.
- Creation Email:** Input field for the email used to create the account.
- Password:** Input field for the user's password.
- Confirm Password:** Input field for the user to confirm their password.
- Country of Residence:** A dropdown menu currently set to "United States of America".
- Work Zip/Postal Code:** Input field for the user's work zip code.
- Occupation:** A dropdown menu currently set to "Physic".

Below the form, there is a "Join" button. A small disclaimer at the bottom of the form reads: "By clicking Join, you agree to our privacy policy and Terms of Use and are NOT a member of our Sites (for now)."

DMA TRAINING
LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
 See slide 65 in DMA Packette

Mental Illness Anxiety Disorders	Inhalers Allergy / Asthma	Hyperlipidemia Statins	Diabetes Endocrine & Metabolic
Abilify Ativan Clozaril Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa	Advair Discus Atrovent Flonase Flovent Loratadine Proventil	Crestor Lipitor Zocor	Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid
Seizures	Gastrointestinal Disorder Constipation	Blood Pressure Meds	Pain & Inflammation
Dilantin Keppra Neurontin Topamax	Colace Miralax Prilosec Protonix Zantac	HCTZ (hydrochlorothiazide) Lisinopril Toprol Tenormin	Flexeril Motrin Norco Tylenol with Codeine Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 1014

DMA Code #2 1965

DMA Code #3 2013

DRUG NAME	Zyprexa
GENERIC NAME	Olanzapine
DOSAGE RANGE	5-20 mg daily
HOW TO TAKE	Orally
USES	To treat mental (moods): Schizophrenia, and / or Bipolar dis.
SIDE EFFECTS	Drowsy, dizzy, dry mouth, weight gain, upset stomach, constipation
WARNINGS	Allergic reaction, dizzy or drowsy (may pass out) Alcohol or marijuana can make you dizzy or drowsy

DRUG NAME	Proventil
GENERIC NAME	Albuterol
DOSAGE RANGE	0.15-2.5 mg/dose Every
HOW TO TAKE	2 puffs 15-30 min before exercise, 4-6 hours as needed
USES	Orally (By mouth/inhale) for Asthma / Allergies
SIDE EFFECTS	Nervousness, shaking, headache, mouth dryness, nausea, dizzy
WARNINGS	May raise blood pressure, chest pain, irregular heart beat, rapid breathing, or confusion; Allergic reaction; Alcohol and weed can cause dizziness

DRUG NAME	ZOCOR
GENERIC NAME	Simvastatin
DOSAGE RANGE	5-40 mg a day
HOW TO TAKE	Orally, once daily in the evening
USES	Help lower bad cholesterol and fats and raise good cholesterol in the blood
SIDE EFFECTS	Mild memory problems or confusion
WARNINGS	May cause OR worsen diabetes; muscle problems; liver problems; allergic reactions

DRUG NAME	Byetta
GENERIC NAME	Exenatide
DOSAGE RANGE	5mcg to 10mcg
HOW TO TAKE	Inject under flesh skin of upper leg, stomach, or upper arm
USES	Treat type 2 diabetes Mellitus
SIDE EFFECTS	Nausea, vomiting, diarrhea, nervous, upset stomach, weight loss
WARNINGS	Blurred vision, drowsiness so don't drive or use machinery or do any activity that you need alertness or clear vision until you can safely

DRUG NAME	Lantus
GENERIC NAME	Insulin Glargine
DOSAGE RANGE	0.2-0.4 units/kg
HOW TO TAKE	Poke finger and take once a day same time
USES	Treat type 1/2 diabetes (controls high blood sugar)
SIDE EFFECTS	Pain, redness, irritation, weight gain
WARNINGS	Do not use when have low blood sugar

can cause blurred vision or dizzy so don't drive, use Machinery, or any activity that require alertness

DRUG NAME	Neurontin
GENERIC NAME	Gabapentin
DOSAGE RANGE	1000mg up to 2400 mg / day
HOW TO TAKE	Orally
USES	Manage epileptic seizures, relieves pain
SIDE EFFECTS	Dizzy, Drowsy, tiredness, loss of coordination, blurred vision, unusual eye movement
WARNINGS	Headache Dizziness and drowsy + blurred vision shaking

DRUG NAME	Zantac
GENERIC NAME	Famotidine
DOSAGE RANGE	No more than 300 mg / day
HOW TO TAKE	Orally every 12 hours
USES	Treat heartburn, acid indigestion, GERD, and gastric ulcers
SIDE EFFECTS	Headache, constipation, Diarrhea
WARNINGS	Allergies, heartburn with lightheadedness/sweating/dizzy Chest / jaw / arm / shoulder pain, bloody vomit

DRUG NAME	Toprol
GENERIC NAME	Metoprolol Succinate
DOSAGE RANGE	50-100 mg / day
HOW TO TAKE	Orally 1/2 a day with meals or after a meal
USES	Chest pain, heart failure, high blood pressure
SIDE EFFECTS	Drowsy, dizzy, tiredness, diarrhea, slow heartbeat
WARNINGS	Allergies, Alcohol or wine can make you more dizzy Do not operate a car or machinery; It can harm baby if pregnant

DRUG NAME	Motrin
GENERIC NAME	Ibuprofen
DOSAGE RANGE	1200-3200 mg / day
HOW TO TAKE	Orally every 4-6 hours with full glass of water
USES	Headache, cold, backache, toothache, mens. cramps, ^{minor} arthritis
SIDE EFFECTS	upset stomach, nausea, vomiting, headache, diarrhea, ^{consti} pain
WARNINGS	Kidney problems, Dehydration, no drugs or IHI ^{or} ^{sun} ^{screen} Stomach bleeding

DRUG NAME	Tylenol with Codeine
GENERIC NAME	Acetaminophen - codeine
DOSAGE RANGE	15- 100 mg / day
HOW TO TAKE	Orally every 4 hours as needed
USES	relieve mild to moderate pain and reduce fever
SIDE EFFECTS	Nausea, vomiting, constipation, drowsy, lighthead
WARNINGS	can cause serious liver damage; No other drugs

slow breathing and severe drowsiness