

Lisa Moore  
5/2/22

DRUG NAME	Abilify
GENERIC NAME	Aripiprazole
DOSAGE RANGE	2mg - 30mg
HOW TO TAKE	x2 day with or without food orally
USES	treats mental/mood, bipolar, Schizophrenia, turrets
SIDE EFFECTS	weight gain, nausea, constipation, insomnia
WARNINGS	lightheadedness, drowsiness, blurred vision, vomiting

DRUG NAME	Flovent
GENERIC NAME	Fluticasone propionate
DOSAGE RANGE	44 mcg - 220 mcg per actuation
HOW TO TAKE	inhaled
USES	to prevent asthma attacks
SIDE EFFECTS	throat irritation, hoarseness, headache
WARNINGS	weakness, tired feeling, or other breathing problems

DRUG NAME	Lipitor
GENERIC NAME	atorvastatin
DOSAGE RANGE	<del>40mg - 80mg</del> 10mg - 80mg
HOW TO TAKE	orally - with or without food
USES	lower blood levels or bad cholesterol
SIDE EFFECTS	joint pain, stuffy nose, sore throat,
WARNINGS	pain burning whe you urinate, weakness

DRUG NAME	Lantus
GENERIC NAME	insulin glargine
DOSAGE RANGE	<del>Self-injectable</del> U-100 pen - 100 units per ml
HOW TO TAKE	Self-injectable
USES	to treat Adults with type 2 diabetes
SIDE EFFECTS	headache, hunger, tremors
WARNINGS	Fast heart rate, Slurred speech, anxiety

DRUG NAME	Keppra
GENERIC NAME	levetiracetam
DOSAGE RANGE	250mg - 100mg
HOW TO TAKE	orally
USES	§ treats symptoms of partial onset seizures
SIDE EFFECTS	dizziness, weakness, tiredness
WARNINGS	changes in mood or behavior, problems with <del>coordination</del> coordination

DRUG NAME	Zantac
GENERIC NAME	Famotidine
DOSAGE RANGE	150 mg - 300 mg
HOW TO TAKE	orally
USES	preventing and relieving heartburn
SIDE EFFECTS	Stomach pain, constipation, Diarrhea, Headache
WARNINGS	May increase your risk of developing pneumonia

DRUG NAME	Toprol
GENERIC NAME	Metoprolol Succinate
DOSAGE RANGE	25 mg - 200 mg
HOW TO TAKE	orally
USES	To treat chest pain, heart failure and high blood pressure
SIDE EFFECTS	dizziness, depression, nightmares, confusion
WARNINGS	Severe dizziness, shortness of breath, fainting

DRUG NAME	Norco
GENERIC NAME	hydrocodone/acetaminophen
DOSAGE RANGE	5 mg - 325 mg
HOW TO TAKE	orally
USES	relieve moderate to severe pain
SIDE EFFECTS	nausea, vomiting, constipation, lightheadedness
WARNINGS	dizziness, or drowsiness

DRUG NAME	Ultram / brand discontinued in the US
GENERIC NAME	tramadol
DOSAGE RANGE	50 mg - 300 mg
HOW TO TAKE	orally
USES	to relieve moderate to severe pain
SIDE EFFECTS	constipation, discouragement, drowsiness
WARNINGS	risk for abuse and addiction can lead to overdose

DRUG NAME	ZOCOR
GENERIC NAME	Simvastatin
DOSAGE RANGE	5 mg - 80 mg
HOW TO TAKE	orally
USES	used to lower levels of bad cholesterol
SIDE EFFECTS	heartburn, gas, bloating, nausea
WARNINGS	loss of appetite, dark urine, yellowing of the skin