

10/30/21
Elizabeth Massing

DMA TRAINING

LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
See slide 65 in DMA Packette

Mental Illness Anxiety Disorders	Inhalers Allergy / Asthma	Hyperlipidemia Statins	Diabetes Endocrine & Metabolic
Abilify Ativan Clozaril Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa	Advair Discus Atrovent Flonase Flovent Loratadine Proventil	Crestor Lipitor Zocor	Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid
Seizures	Gastrointestinal Disorder Constipation	Blood Pressure Meds	Pain & Inflammation
Dilantin Keppra Neurontin Topamax	Colace Miralax Prilosec Protonix Zantac	HCTZ (hydrochlorothiazide) Lisinopril Toprol Tenormin	Flexeril Motrin Norco Tylenol with Codeine Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 1737 DMA Code #2 1978 DMA Code #3 9532

DRUG NAME	KEPTIV
GENERIC NAME	levetiracetam
DOSAGE RANGE	TAB: 250mg, 500mg, 750mg, 1000mg; SOL: 100mg Per ML
HOW TO TAKE	500mg - 1500mg PO/IV q12h
USES	Partial Seizures, Primary generalized
SIDE EFFECTS	vertigo, fatigue, headache, irritability, vomiting
WARNINGS	PSYCHIATRIC disorders,

DRUG NAME	Protonix
GENERIC NAME	Pantoprazole
DOSAGE RANGE	TAB: 20mg, 40mg; DR Granule 40mg Per PKT
HOW TO TAKE	20mg PO/qd x 4-8 wk
USES	GERD, hypersecretory conditions, UPPER GI bleeding
SIDE EFFECTS	fatigue, hypomagnesemia, vitamin B12 deficiency
WARNINGS	long term use, high dose vs

DRUG NAME	Toprol
GENERIC NAME	metoprolol succinate
DOSAGE RANGE	TABER 25mg, 50mg, 100mg, 200mg
HOW TO TAKE	50-200mg PO qd
USES	HTN, angina, heart failure - associated risk reduction
SIDE EFFECTS	Fatigue, dizziness, diarrhea, PROSTITIS
WARNINGS	Thyroid disorder, alcohol use, major surgery

DRUG NAME	VIIAM Norco
GENERIC NAME	hydrocodone hydrocodone/acetaminophen
DOSAGE RANGE	5mg/325mg, 7.5mg/325mg, 10mg/325mg
HOW TO TAKE	2.5-10mg hydrocodone PO q4-6h PRN
USES	Pain moderate-severe
SIDE EFFECTS	constipation, light-headedness, drowsiness
WARNINGS	GI obstruction, head injury, hepatic impairment

DRUG NAME	Depakote
GENERIC NAME	Divalproex Sodium
DOSAGE RANGE	DB TAB 125mg, 250mg, 500mg
HOW TO TAKE	250-500mg PO BID
USES	absence seizures, bipolar disorder manic acute
SIDE EFFECTS	headache, nausea, vomiting, asthenia, somnolence
WARNINGS	bleeding risk, hepatic dz, renal impairment

DRUG NAME	1110741
GENERIC NAME	Oxcarbazepine
DOSAGE RANGE	TAB: 150mg, 300mg, 600mg; SUSP: 300mg Per 5mL
HOW TO TAKE	600 by mouth (PO) Twice daily (BID)
USES	Seizures, bipolar disorder, Trigeminal neuralgia
SIDE EFFECTS	Dizziness, Coordination problems, Nausea, Double vision
WARNINGS	Do not drink alcohol, Avoid becoming dehydrated.

DRUG NAME	Advair Diskus
GENERIC NAME	Fluticasone Propionate / Salmeterol Inhaled
DOSAGE RANGE	100 mcg / 50 mcg per blister, 250 mcg / 50 mcg per blister, 500mcg
HOW TO TAKE	1 puff Inhaled q12h.
USES	asthma, COPD maintenance
SIDE EFFECTS	Cold symptoms, Thrush, headache, muscle pain, bone pain, back pain
WARNINGS	Do not use if severe allergy to milk protein

DRUG NAME	Ativan
GENERIC NAME	Lorazepam
DOSAGE RANGE	TAB: 0.5mg, 1mg, 2mg; INJ: 2mg per mL, 4mg per mL
HOW TO TAKE	2-3mg/day PO/IM/IV divided BID-TID, max 10mg/Per Day
USES	Anxiety, insomnia, Short term, Status epilepticus
SIDE EFFECTS	Dizziness, Drowsiness, weakness, feeling unsteady.
WARNINGS	Avoid alcohol, operating heavy machinery.

DRUG NAME	ZOCOR
GENERIC NAME	Simvastatin
DOSAGE RANGE	TAB: 10mg, 20mg, 40mg, 80mg
HOW TO TAKE	20-40 mg PO qd Max: 40mg/Day
USES	Primary hyperlipidemia, hypertriglyceridemia
SIDE EFFECTS	headache, stomach pain, constipation, cold symptoms.
WARNINGS	Grapefruit may interact with Simvastatin and lead to unwanted side effects.

DRUG NAME	AP: dfg
GENERIC NAME	INSULIN glulisine
DOSAGE RANGE	INJ (SoloStar U-100 Pen) 100 units per mL; INJ: (U-100 Vial) 100 units
HOW TO TAKE	Diabetes mellitus
USES	individualize dose SC/IV BID-QID
SIDE EFFECTS	localized cutaneous amyloidosis, renal impairment
WARNINGS	hypersensitivity to metacresol