DMA TRAINING

LIST OF MEDICATIONS TO COMPLETE FOR DMA TRAINING

Use the attached forms to look up each of the medications listed below. Each line must be completed and turned in the day that you do your final DMA Evaluation with your ROM (Regional Operation Manager) for your area. You will not be able to become DMA certified until all of the forms are completed [48]
See slide 65 in DMA Packette

Mental Illness **Anxiety Disorders**	**Inhalers** **Allergy / Asthma**	**Hyperlipidemia** **Statins**	**Diabetes** **Endocrine & Metabolic**
Abilify Ativan Clozaril Depakote Haldol Invega Klonopin Lamictal Lithium Risperdal Seroquel Tripleptal Zyprexa	Advair Discus Atrovent Flonase Flovent Loratadine Proventil	Crestor Lipitor Zocor	Apidra Byetta Glucophage Glyburide Lantus Levemir Levothyroxine Novolog Synthroid
Seizures	**Gastrointestinal Disorder** **Constipation**	**Blood Pressure Meds**	**Pain & Inflammation**
Dilantin Keppra Neurontin Topamax	Colace Miralax Prilosec Protonix Zantac	HCTZ (hydorchlorothiazide) Lisinopril Toprol Tenormin	Flexeril Motrin Norco Tylenol with Codeine Ultram

App/Website: Epocrates

Please complete 10 of the above medications prior to attending DMA class.

DMA Code #1 _____ DMA Code #2 _____ DMA Code #3 _____

DRUG NAME	Norco
GENERIC NAME	hydrocodone / acetaminophen
DOSAGE RANGE	5mg / 325mg ; 7.5mg / 325mg, 10mg / 325mg
HOW TO TAKE	by mouth (PO)
USES	Pain
SIDE EFFECTS	nausea, sedation, vomiting, constipation, psychomotor impairment
WARNINGS	overdose, death, addiction, Abuse, REMS, Resp. Depression, accidental Ingestion by children can be fatal, Hepatotoxicity.

DRUG NAME	Tylenol with Codeine
GENERIC NAME	
DOSAGE RANGE	
HOW TO TAKE	~~Discontinued in US~~
USES	
SIDE EFFECTS	See Back
WARNINGS	

DRUG NAME	Toprol-XL
GENERIC NAME	metoprolol succinate
DOSAGE RANGE	12.5mg - 400mg
HOW TO TAKE	by mouth (PO)
USES	Antianginals, Beta Blockers, Headache / Migraine
SIDE EFFECTS	nightmares, bradycardia, rash, fatigue, dizziness, dyspnea
WARNINGS	

DRUG NAME	Lipitor
GENERIC NAME	atorvastin
DOSAGE RANGE	10mg - 80mg
HOW TO TAKE	by mouth (PO)
USES	Dyslipidemia: HMG-CoA Reductase Inhibitors (Statin
SIDE EFFECTS	UTI, musculoskeletal pain, extremity pain, cognitive impairme
WARNINGS	No Black Box Warnings

DRUG NAME	Haldol
GENERIC NAME	haloperidol
DOSAGE RANGE	0.5mg - 20mg
HOW TO TAKE	by mouth (PO)
USES	Antipsychotics, 1st Gen. Tic Disorders / Tourette Syndrom
SIDE EFFECTS	lethargy, anxiety, impaired body temp. regulation, insomnia
WARNINGS	Dementia Related Psychosis

Tylenol w/ Codeine
acetaminophen / codeine
TAB 300mg/15mg, 300mg/30mg, 300mg/60mg.
SOL 120mg/12mg per 5mL
· Elixir, Tablet, Solution - By mouth (PO)
· Pain, cough
nausea, abdominal pain, rash, euphoria, sedation
accidental overdose, death, addiction, Abuse, REMS,
Resp. Depression, ultra-Rapid Metabolism of
Codeine + other Res. Depression in children,
· Neonatal opioid withdrawal syndrome,
Hepatotoxicity, Accidental ingestion of
children can cause death.

DRUG NAME	Ativan
GENERIC NAME	Lorazepam
DOSAGE RANGE	0.5, 1mg, 2mg, INJ: 2mg per mL, 4mg per ML
HOW TO TAKE	by mouth (PO), intramuscular (IM) Intravenous (IV)
USES	Antiemetics, "benzos", Alcohol Dependence, Insomnia, Sedation, Seizures
SIDE EFFECTS	sedation, constipation, menstural irregularities, dizziness, ataxia
WARNINGS	Addiction, Dependence, profound sedation, resp. depression, coma, death.

DRUG NAME	Clozaril
GENERIC NAME	Clozapine
DOSAGE RANGE	25mg, 100 mg
HOW TO TAKE	by mouth
USES	Antipsychotics (schizophrenia, psychosis, depression, A12
SIDE EFFECTS	weight gain, tachycardia, fever, tremor, visual disturbance
WARNINGS	Neutropenia, seizures, myocarditis, Dementia related Psychosis cardiomyopathy, orthostatic hypotension.

DRUG NAME	Klonopin
GENERIC NAME	Clonazepam
DOSAGE RANGE	0.5mg, 1mg, 2mg
HOW TO TAKE	by mouth
USES	seizure disorders, Tourette syndrome, "Benzos", sleep related movement
SIDE EFFECTS	depression, incontinence, rash, confusion, amnesia, drowsiness
WARNINGS	Addiction, Abuse, profound resp. depression, coma, death, Dependence.

DRUG NAME	Levothyroxine
GENERIC NAME	Euthyrox, Levothroid, Levoxyl, Synthroid, Thyquidity, Tirosint
DOSAGE RANGE	13 mcg, 25, 50, 75, 88, 100, 112, 125, 137, 150, 175, 200, (Cap)
HOW TO TAKE	by mouth (PO) intravenous (IV) intramuscular (IM)
USES	Thyroid Disorders: Hypothyroidism
SIDE EFFECTS	heat intolerance, insomnia, anxiety, weight loss, palpitations
WARNINGS	Not for weight loss, life-threatening toxicity.

Eutirox

DRUG NAME	Depakote
GENERIC NAME	Divalproex sodium
DOSAGE RANGE	30mg - 500mg
HOW TO TAKE	By mouth (PO)
USES	Bi Polar disorder, Headache/Migraines, Seizure Disorders
SIDE EFFECTS	blurred vision, photosensitivity, vomiting, headache, diarrhea.
WARNINGS	Hepatotoxicity, Fetal Risk, Pancreatitis

Levothyroxine dosage Range

Tab: 25, 50, 75, 88, 100, 112, 125, 137, 175, 200, 300 mcg.

INJ: Varies