

\_H\_\_ Thrill

**F.** Turbulent sounds occurring between normal heart sounds. Characterized by loudness, pitch, shape, duration. Graded I-VI

\_L\_ 3<sup>rd</sup> Heart Sound

**D.** Small red streaks under fingernails

\_I\_ Troponins

**G.** HR > 100

\_N\_\_ HDL

**E.** HR < 60

\_F\_\_ Heart murmurs

**B.** The amount of blood pumped by each ventricle in 1 min. Calculated by Stroke Volume X HR

\_D\_\_ Splinter hemorrhages

**M.** Peripheral resistance against which the left ventricle must pump

\_G\_\_ Tachycardia

**A.** Turbulent flow through artery caused by aneurysm or arterial obstruction

\_E\_\_ Bradycardia

**I.** Contractile proteins released after MI. Both T and I samples test for cardiac injury/ischemia

\_A\_\_ Arterial bruit

**L.** Extra heart sound heard in EARLY diastole, similar to the sound of a gallop. Causes include Left Ventricular failure, volume overload, aortic or tricuspid regurgitation

\_J\_\_ LDL

**K.** Alteration of spine often in elderly resulting in altered chest landmarks, auscultation, and percussion

\_B\_\_ Cardiac Output

**C.** The volume of blood stretching the ventricles at the end of diastole before the next contraction

\_K\_\_ Kyphosis

**H.** Vibration of vessel or chest

\_C\_\_ Preload

**N.** Lipoprotein considered to be the "good cholesterol"

\_M\_\_ Afterload

**J.** Lipoprotein considered to be "lethal cholesterol"