

## Cardiac Assessment Definition Match

\_\_H\_\_ Thrill

F. Turbulent sounds occurring between normal heart sounds. Characterized by loudness, pitch, shape, duration. Graded I-VI

\_\_L\_\_ 3<sup>rd</sup> Heart Sound

D. Small red streaks under fingernails

\_\_I\_\_ Troponins

G. HR > 100

\_\_N\_\_ HDL

E. HR < 60

\_\_F\_\_ Heart murmurs

B. The amount of blood pumped by each ventricle in 1 min. Calculated by Stroke Volume X HR

\_\_D\_\_ Splinter hemorrhages

M. Peripheral resistance against which the left ventricle must pump

\_\_G\_\_ Tachycardia

A. Turbulent flow through artery caused by aneurysm or arterial obstruction

\_\_E\_\_ Bradycardia

I. Contractile proteins released after MI. Both T and I samples test for cardiac injury/ischemia

\_\_A\_\_ Arterial bruit

L. Extra heart sound heard in EARLY diastole, similar to the sound of a gallop. Causes include Left Ventricular failure, volume overload, aortic or tricuspid regurgitation

\_\_J\_\_ LDL

K. Alteration of spine often in elderly resulting in altered chest landmarks, auscultation, and percussion

\_\_B\_\_ Cardiac Output

C. The volume of blood stretching the ventricles at the end of diastole before the next contraction

\_\_K\_\_ Kyphosis

H. Vibration of vessel or chest

\_\_C\_\_ Preload

N. Lipoprotein considered to be the "good cholesterol"

\_\_M\_\_ Afterload

J. Lipoprotein considered to be "lethal cholesterol"