

Cardiac Assessment Definition Match

__**H**__ Thrill

__**L**__ 3rd Heart Sound

__**I**__ Troponins

__**N**__ HDL

__**F**__ Heart murmurs

__**D**__ Splinter hemorrhages

__**G**__ Tachycardia

__**E**__ Bradycardia

__**A**__ Arterial bruit

__**J**__ LDL

__**B**__ Cardiac Output

__**K**__ Kyphosis

__**C**__ Preload

__**M**__ Afterload

F. Turbulent sounds occurring between normal heart sounds. Characterized by loudness, pitch, shape, duration. Graded I-VI

D. Small red streaks under fingernails

G. HR > 100

E. HR < 60

B. The amount of blood pumped by each ventricle in 1 min. Calculated by Stroke Volume X HR

M. Peripheral resistance against which the left ventricle must pump

A. Turbulent flow through artery caused by aneurysm or arterial obstruction

I. Contractile proteins released after MI. Both T and I samples test for cardiac injury/ischemia

L. Extra heart sound heard in EARLY diastole, similar to the sound of a gallop. Causes include Left Ventricular failure, volume overload, aortic or tricuspid regurgitation

K. Alteration of spine often in elderly resulting in altered chest landmarks, auscultation, and percussion

C. The volume of blood stretching the ventricles at the end of diastole before the next contraction

H. Vibration of vessel or chest

N. Lipoprotein considered to be the “good cholesterol”

J. Lipoprotein considered to be “lethal cholesterol”