

Additional IV medication information:

**Morphine:**

May give undiluted, or w sterile water/0.9 NaCl 1-2 mg/ml for IV injection. For continuous infusion, dilute to concentration of 0.1-1mg/mL in dextrose. Always admin very slowly, store at room temperature.

Incompatible: Amphotericin B complex, cefepime, doxorubicin, phenytoin

Compatible: Amiodarone, atropine, bumetanide, bupivacaine, dexmedetomidine, diltiazem, diphenhydramine, dobutamine, dopamine, glycopyrrolate, heparin, hydroxyzine, lidocaine, lorazepam, magnesium, midazolam, milrinone, nitroglycerin, potassium, propofol

**Infliximab:**

Reconstitute each vial w 10 mL sterile water for injection using 21 gauge or smaller needle, direct stream of sterile water for injection to glass wall of vial. Swirl vial gently to dissolve contents. Allow solution to stand for 5 min, inject into 250 ml bag 0.9 NaCL, gently mix, concentration 0/4- 4 mg/ml, begin infusion w/in 3 hr after reconstitution. Admin infusion over at least 2 hrs using a low protein-binding filter. Refrigerate vials, solution should appear colorless to light yellow and opalescent, do not use if colored or if particulate forms.

Incompatibilities: Do not infuse in same IV line as other agents