

Module Report

Tutorial: Real Life RN Medical Surgical 4.0

Module: GI Bleed



Individual Name: Ryan Clagett

Institution: Margaret H Rollins SON at Beebe Medical Center

Program Type: Diploma

Standard Use Time and Score

	Date/Time	Time Use	Score
GI Bleed	3/5/2024 11:11:57 AM	25 min	Satisfactory

Reasoning Scenario Details

GI Bleed - Use on 3/5/2024 10:46:41 AM

Reasoning Scenario Performance Related to Outcomes:

*See Score Explanation and Interpretation below for additional details.

Body Function	Strong	Satisfactory	Needs Improvement
Cardiac Output and Tissue Perfusion	100%		
Cognition and Sensation	50%		50%
Ingestion, Digestion, Absorption & Elimination	100%		
Regulation and Metabolism	100%		

NCLEX RN	Strong	Satisfactory	Needs Improvement
RN Management of Care	100%		
RN Health Promotion and Maintenance	50%		50%
RN Psychosocial Integrity	100%		
RN Pharmacological and Parenteral Therapies	100%		
RN Reduction of Risk Potential	100%		
RN Physiological Adaptation	100%		

QSEN	Strong	Satisfactory	Needs Improvement
Safety	100%		
Patient-Centered Care	83.3%		16.7%
Evidence Based Practice	100%		

Decision Log:

Optimal Decision	
Scenario	Nurse Esther listens to bowel sounds.
Question	Nurse Esther listens to Ms. Lieberman's abdomen in all four quadrants and determines Ms. Lieberman's bowel sounds are hyperactive. Listen to the four audio clips. Which of the following sounds is an expected finding for Ms. Lieberman?
Selected Option	Option C: Audio clip of bowel sounds occurring 45 times in 1 min.
Rationale	Bowel sounds are clicks and gurgles heard in the abdomen. Bowel sounds within the expected reference range are irregular sounds that occur five to 35 times a minute. This finding indicates hyperactive bowel sounds. Therefore, this is the expected finding for this client.

Optimal Decision	
Scenario	Ms. Lieberman reports she feels lightheaded and dizzy.
Question	Ms. Lieberman states she is feeling lightheaded and dizzy. Her skin color is pale. Which of the following should be Nurse Esther's priority action?
Selected Option	Measure Ms. Lieberman's vital signs.
Rationale	The client is at risk for hypovolemic shock due to the loss of extracellular fluid and blood. Clinical manifestations of hypovolemic shock include hypotension and tachycardia. Therefore, the nurse should assess the client's status by obtaining her vital signs.

Optimal Decision	
Scenario	Nurse Esther obtains Ms. Lieberman's vital signs after she reports feeling faint.
Question	Ms. Lieberman reports feeling worse and her vital signs are: BP 94/56 mm Hg, pulse 110/min, respirations 26/min, and SaO2 94%. Nurse Esther starts oxygen at 2 L/min. Which of the following should be Nurse Esther's priority action?
Selected Option	Lower the head of the bed.
Rationale	Using the ABC priority-setting framework, the priority response is to promote improved circulation by lowering the head of the bed and elevating the client's feet. This action can prevent hypovolemic shock until adequate blood volume is restored.

Optimal Decision

Scenario	The unit of packed RBCs for Ms. Lieberman arrives on the unit.
Question	Nurse Esther is preparing to administer a unit of packed RBCs to Ms. Lieberman. Which of the following actions should Nurse Esther perform prior to administering the blood?
Selected Option	Ask Ms. Lieberman if she has experienced a reaction with any previous blood transfusions.
Rationale	A transfusion reaction can be caused by the development of antibodies to the donor leukocytes. This reaction is more likely to occur when a client has had blood transfusions before, as well as a history of prior blood transfusion reactions. Therefore, this is the appropriate action for the nurse to take.

Optimal Decision	
Scenario	Nurse Esther is ready to administer the first unit of packed RBCs.
Question	Identify the correct sequence of actions for blood administration after Nurse Esther performs hand hygiene and applies gloves. (Reorder the steps by dragging them into the desired sequence.)
Selected Ordering	Spike and prime the Y-set tubing with the 0.9% sodium chloride solution. Attach the tubing to the IV catheter and begin to infuse the 0.9% sodium chloride solution. Gently rotate the bag of packed RBCs. Attach the packed RBCs bag to the Y-set tubing. Turn off the 0.9% sodium chloride solution. Begin to infuse the packed RBCs.
Rationale	The first action the nurse should do is insert one of the spikes of the Y-set into the 0.9% sodium chloride solution bag, prime the tubing with the 0.9% sodium chloride solution, and start slowly infusing the solution into the client's IV. Next, the nurse should gently rotate the bag to mix the blood cells with the plasma. Then, the nurse should spike the blood bag with the remaining spike on the Y-set tubing and turn off the 0.9% sodium chloride solution by closing the clamp. Lastly, the nurse needs to open the clamp to allow the blood to infuse.

Optimal Decision	
Scenario	Ms. Lieberman is restless, her face is flushed, and she reports having a headache.
Question	Nurse Esther notes Ms. Lieberman is restless, her face is flushed, and she reports having a headache. Her vital signs include: temperature 38.8° C (101.8° F), pulse 96/min, respirations 22/min, and BP 103/60 mm Hg. Which of the following is an appropriate action for Nurse Esther to take?
Selected Option	Stop the blood transfusion.
Rationale	In the presence of a febrile reaction, the client's blood is sensitive to some component of the donor's blood. To prevent further exposure to the sensitizing component, the transfusion should be stopped immediately.

Scenario	Using an SBAR format, write the information Nurse Esther should give to Ms. Lieberman's provider when calling about her response to the blood. Refer to the EMR documents for needed information.
-----------------	---

Question	Using an SBAR format, write the information Nurse Esther should give to Ms. Lieberman's provider when calling about her response to the blood. Refer to the EMR documents for needed information. (Enter your response, then click on the submit button at the bottom of the screen. Compare your response to the one provided.)
Selected Option	S - JL was admitted for abd cramping pain, monitoring of stool, and receiving a transfusion of PRBCs. B - Administered infusion of PRBCs at 1545. A - 15 mins after the infusion, JL complained of feeling "chilly" and "aching all over." Transfusion stopped promptly. T 38.8 C, RR 22, BP 103/60. R - Recommend prescribing antipyretic and continuing IVF NS to KVO.
Rationale	The following information should be shared with Ms. Lieberman's provider when calling about her response to the blood. Situation: Dr. McGuire, this is Esther - RN. I am taking care of Ms. Lieberman in room 5206. She is a 36-year-old client admitted from the ED today for a GI bleed. She's had one unit of packed RBCs and part of the second unit of blood. I stopped the second unit because I believe she is having a transfusion reaction. Her baseline temperature was 98.6 and is now 101.8. Ms. Lieberman reports having a headache, chills, and is restless. She does not have any evidence of a rash at this time. Background: Ms. Lieberman has a history of Crohn's disease and intermittent gastritis. Six months ago she had an ileostomy and started on infliximab IV every 8 weeks. Her last infusion was 7 weeks ago. Assessment: Her hemoglobin was 7 g/dL and her hematocrit was 21% in the ED. When she arrived to the medical surgical unit, her BP was 94/56 and her pulse 110, but now her BP is 110/70 and her pulse is 110. At this time, her respirations are 26, her SaO2 is 97%, and her temperature is 101.8. I have discontinued the second unit of blood and plan to send both the bags of blood to the lab per protocol. I hung a new bag of 0.9% sodium chloride to keep the line open. Recommendations: Ms. Lieberman is requesting ibuprofen for her headache, which would also bring her fever down. Could I have an prescription for an antipyretic, and do you want to continue the IV infusion of 0.9% sodium chloride at 150 mL/hr?

Optimal Decision	
Scenario	Dr. March tells Ms. Lieberman that he recommends an endoscopy. Ms. Lieberman is informed about the procedure. She agrees to the procedure and signs the consent form.
Question	Nurse Esther is reinforcing teaching with Ms. Lieberman, who is scheduled for an endoscopy in the morning. Which of the following should Nurse Esther include in the teaching?
Selected Option	"A medication to reduce oral secretions may be administered."
Rationale	The nurse could administer atropine (Sal-Tropine), a muscarinic antagonist, to inhibit salivary and bronchial secretions.

Optimal Decision	
Scenario	Nurse Esther is calculating the number of milliliters of morphine sulfate to administer.
Question	Nurse Esther is preparing to administer morphine 4 mg IV. Available is morphine 8 mg/mL. How many mL should the nurse administer? (Round the answer to the nearest tenth.).

Selected Option	0.5
Rationale	Follow these steps for the Ratio and Proportion method of calculation: Step 1: What is the unit of measurement the nurse should calculate? mL Step 2: What is the dose the nurse should administer? Dose to administer = Desired 4 mg Step 3: What is the dose available? Dose available = Have 8 mg Step 4: Should the nurse convert the units of measurement? No Step 5: What is the quantity of the dose available? 1 mL Step 6: Set up an equation and solve for X. $\text{Have} \times \text{Desired} = \text{Quantity} \times X$ $8 \text{ mg} \times 1 \text{ mL} = 4 \text{ mg} \times X$ $X = 0.5 \text{ mL}$ Step 7: Round if necessary. Step 8: Determine whether the amount to administer makes sense. If there are 8 mg/mL and the prescription reads 4 mg, it makes sense to administer 0.5 mL. The nurse should administer morphine 0.5 mL IV. Follow these steps for the Desired Over Have method of calculation: Step 1: What is the unit of measurement the nurse should calculate? mL Step 2: What is the dose the nurse should administer? Dose to administer = Desired 4 mg Step 3: What is the dose available? Dose available = Have 8 mg Step 4: Should the nurse convert the units of measurement? No Step 5: What is the quantity of the dose available? 1 mL Step 6: Set up an equation and solve for X. $\text{Desired} \times \text{Quantity} = \text{Have} \times X$ $4 \text{ mg} \times 1 \text{ mL} = 8 \text{ mg} \times X$ $X = 0.5 \text{ mL}$ Step 7: Round if necessary. Step 8: Determine whether the amount to administer makes sense. If there are 8 mg/mL and the prescription reads 4 mg, it makes sense to administer 0.5 mL. The nurse should administer morphine 0.5 mL IV. Follow these steps for the Dimensional Analysis method of calculation: Step 1: What is the unit of measurement the nurse should calculate? (Place the unit of measure being calculated on the left side of the equation.) $X \text{ mL} =$ Step 2: Determine the ratio that contains the same unit as the unit being calculated. (Place the ratio on the right side of the equation, ensuring that the unit in the numerator matches the unit being calculated.) $1 \text{ mL} \times \frac{8 \text{ mg}}{4 \text{ mg}} = X$ Step 3: Place any remaining ratios that are relevant to the item on the right side of the equation, along with any needed conversion factors, to cancel out unwanted units of measurement. $1 \text{ mL} \times \frac{8 \text{ mg}}{4 \text{ mg}} = X$ Step 4: Solve for X. $X = 0.5 \text{ mL}$ Step 5: Round if necessary. Step 6: Determine whether the amount to administer makes sense. If there are 8 mg/mL and the prescription reads 4 mg, it makes sense to administer 0.5 mL. The nurse should administer morphine 0.5 mL IV.

Optimal Decision	
Scenario	Nursing considerations Nurse Esther takes when administering morphine.

Question	Nurse Esther is preparing to administer 4 mg of morphine IV bolus to Ms. Lieberman. Which of the following actions should Nurse Esther take?
Selected Option	Infuse morphine at a rate of 1 mg/min.
Rationale	To prevent serious adverse reactions, such as respiratory arrest, the nurse should inject the medication at a rate of 1 mg/min.

Optimal Decision	
Scenario	Nurse Esther talks to Ms. Lieberman about needing several drinks after work to relax.
Question	Ms. Lieberman tells Nurse Esther she has a stressful job working in the city as a stockbroker, and that sometimes at night she has up to five drinks. Which of the following is an appropriate statement made by Nurse Esther?
Selected Option	"Tell me more about the stress you are feeling."
Rationale	Providing an open-ended statement, along with active listening, allows the client to express her thoughts and feelings. It also establishes trust.

Scenario	Identify five stress management strategies Nurse Esther should recommend to Ms. Lieberman to promote a healthier lifestyle.
Question	Identify five stress management strategies Nurse Esther should recommend to Ms. Lieberman to promote a healthier lifestyle. (Enter your response, then click on the submit button at the bottom of the screen. Compare your response to the one provided.)
Selected Option	1) Exercise, 2) Following a high-protein, high-vitamin diet, 3) Finding a hobby or fun activity to participate in outside of work, 4) Call friends after work to hang out, and 5) Take time to meditate prior to going to sleep.
Rationale	Identify five stress management strategies Nurse Esther should recommend to Ms. Lieberman to promote a healthier lifestyle.1. Perform light, regular exercise.2. Write in a journal.3. Listen to music.4. Consider a pet.5. Get adequate sleep.6. Promote relaxation through use of progressive muscle relaxation, guided imagery, massage therapy, humor, and/or yoga.7. Enhance her social support system, such as an ostomy support group, AA, and/or coping support group.8. Evaluate current job, lifestyle, and home location.

Optimal Decision	
Scenario	Nurse Esther discusses diet with Ms. Lieberman.
Question	Nurse Esther is reinforcing diet teaching with Ms. Lieberman. Which of the following dietary recommendations should she make?
Selected Option	Eat foods that are high protein.
Rationale	Clients who have Crohn's disease are at risk for malnutrition because they may attempt to control symptoms by restricting their diet. Additionally, clients who have Crohn's disease are at risk for malabsorption of nutrients. The client should be instructed to maximize her nutrition by eating foods high in protein.

Scenario	Nurse Esther provides Ms. Lieberman with educational material to take home.
-----------------	---

Question	Nurse Esther provides Ms. Lieberman with information about health promotion. Which of the following should she include in the teaching?
Selected Option	Urge Ms. Lieberman to run 30 min, 3 times a week.
Rationale	Clients who have Crohn's disease are at risk for fluid volume deficit. Strenuous activity, such as running, causes heavy perspiration and fluid loss. Therefore, this type of strenuous activity should be avoided.

Individual Report – Score Explanation and Interpretation

Reasoning Scenario Information:

Reasoning Scenario Information provides the date, time and duration of use, along with the score earned for each attempt. A Reasoning Scenario Performance score of Strong, Satisfactory, or Needs Improvement is provided for each attempt. This information is also provided for the Optimal Decision Mode if it has been enabled.

Reasoning Scenario Performance Scores:

Strong	Exhibits optimal reasoning that results in positive outcomes in the care of clients and resolution of problems.
Satisfactory	Exhibits reasoning that results in mildly helpful or neutral outcomes in the care of clients and resolution of problems.
Needs Improvement	Exhibits reasoning that results in harmful or detrimental outcomes in the care of clients and resolution of problems.

Reasoning Scenario Performance Related to Outcomes:

A clinical reasoning performance score related to each outcome is provided. Outcomes associated with student responses are listed in the report. The number across from each outcome indicates the percentage of responses associated with the level of performance of that outcome.

NCLEX® Client Need Categories:

Management of Care	Providing integrated, cost-effective care to clients by coordinating, supervising, and/or collaborating with members of the multi-disciplinary health care team.
Safety and Infection Control	Incorporating preventative safety measures in the provision of client care that provides for the health and well-being of clients, significant others, and members of the health care team.
Health Promotion and Maintenance	Providing and directing nursing care that encourages prevention and early detection of illness, as well as the promotion of health.
Psychosocial Integrity	Promoting mental, emotional, and social well-being of clients and significant others through the provision of nursing care.
Basic Care and Comfort	Promoting comfort while helping clients perform activities of daily living.
Pharmacological and Parenteral Therapies	Providing and directing administration of medication, including parenteral therapy.
Reduction of Risk Potential	Providing nursing care that decreases the risk of clients developing health-related complications.

Physiological Adaptation	Providing and directing nursing care for clients experiencing physical illness.
---------------------------------	---

Quality and Safety Education for Nurses (QSEN)

Safety	The minimization of risk factors that could cause injury or harm while promoting quality care and maintaining a secure environment for clients, self, and others.
Patient-Centered Care	The provision of caring and compassionate, culturally sensitive care that is based on a client's physiological, psychological, sociological, spiritual, and cultural needs, preferences, and values
Evidence Based Practice	The use of current knowledge from research and other credible sources, upon which clinical judgment and client care are based.
Informatics	The use of information technology as a communication and information gathering tool that supports clinical decision making and scientifically based nursing practice.
Quality Improvement	Care related and organizational processes that involve the development and implementation of a plan to improve health care services and better meet the needs of clients.
Teamwork and Collaboration	The delivery of client care in partnership with multidisciplinary members of the health care team, to achieve continuity of care and positive client outcomes.

Body Function

Cardiac Output and Tissue Perfusion	The anatomical structures (heart, blood vessels, and blood) and body functions that support adequate cardiac output and perfusion of body tissues.
Cognition and Sensation	The anatomical structures (brain, central and peripheral nervous systems, eyes and ears) and body functions that support perception, interpretation, and response to internal and external stimuli.
Excretion	The anatomical structures (kidney, ureters, and bladder) and body functions that support filtration and excretion of liquid wastes, regulate fluid and electrolyte and acid-base balance.
Immunity	The anatomic structures (spleen, thymus, bone marrow, and lymphatic system) and body functions related to inflammation, immunity, and cell growth.
Ingestion, Digestion, Absorption and Elimination	The anatomical structures (mouth, esophagus, stomach, gall bladder, liver, small and large bowel, and rectum) and body functions that support ingestion, digestion, and absorption of food and elimination of solid wastes from the body.
Integument	The anatomical structures (skin, hair, and nails) and body functions related to protecting the inner organs from the external environment and injury.
Mobility	The anatomical structures (bones, joints, and muscles) and body functions that support the body and provide its movement.

Oxygenation	The anatomical structures (nose, pharynx, larynx, trachea, and lungs) and body functions that support adequate oxygenation of tissues and removal of carbon dioxide.
Regulation and Metabolism	The anatomical structures (pituitary, thyroid, parathyroid, pancreas, and adrenal glands) and body functions that regulate the body's internal environment.
Reproduction	The anatomical structures (breasts, ovaries, fallopian tubes, uterus, vagina, vulva, testicles, prostate, scrotum, and penis) and body functions that support reproductive functions.

Decision Log

Information related to each question answered in a scenario attempt is listed in the report. A brief description of the scenario, question, selected option and rationale for that option are provided for each question answered. The words "Optimal Decision" appear next to the question when the most optimal option was selected.

The rationale for each selected option may be used to guide remediation. A variety of learning resources may be used in the review process, including related ATI Review Modules.

Module Report

Tutorial: Real Life RN Medical Surgical 4.0

Module: GI Bleed



Individual Name: Ryan Clagett

Institution: Margaret H Rollins SON at Beebe Medical Center

Program Type: Diploma

Standard Use Time and Score

	Date/Time	Time Use	Score
GI Bleed	3/5/2024 10:45:53 AM	1 hr 20 min	Needs Improvement
This attempt ended prematurely due to a detrimental decision or a series of missteps.			

Reasoning Scenario Details

GI Bleed - Use on 3/5/2024 9:26:28 AM

Reasoning Scenario Performance Related to Outcomes:

*See Score Explanation and Interpretation below for additional details.

Body Function	Strong	Satisfactory	Needs Improvement
Cardiac Output and Tissue Perfusion	60%	40%	
Ingestion, Digestion, Absorption & Elimination	66.7%		33.3%
Regulation and Metabolism			100%

NCLEX RN	Strong	Satisfactory	Needs Improvement
RN Management of Care			100%
RN Pharmacological and Parenteral Therapies	50%		50%
RN Reduction of Risk Potential	100%		
RN Physiological Adaptation	60%	40%	

QSEN	Strong	Satisfactory	Needs Improvement
Safety	75%		25%

Patient-Centered Care	66.7%	33.3%	
Evidence Based Practice		50%	50%

Decision Log:

Optimal Decision	
Scenario	Nurse Esther listens to bowel sounds.
Question	Nurse Esther listens to Ms. Lieberman's abdomen in all four quadrants and determines Ms. Lieberman's bowel sounds are hyperactive. Listen to the four audio clips. Which of the following sounds is an expected finding for Ms. Lieberman?
Selected Option	Option C: Audio clip of bowel sounds occurring 45 times in 1 min.
Rationale	Bowel sounds are clicks and gurgles heard in the abdomen. Bowel sounds within the expected reference range are irregular sounds that occur five to 35 times a minute. This finding indicates hyperactive bowel sounds. Therefore, this is the expected finding for this client.

Optimal Decision	
Scenario	Ms. Lieberman reports she feels lightheaded and dizzy.
Question	Ms. Lieberman states she is feeling lightheaded and dizzy. Her skin color is pale. Which of the following should be Nurse Esther's priority action?
Selected Option	Measure Ms. Lieberman's vital signs.
Rationale	The client is at risk for hypovolemic shock due to the loss of extracellular fluid and blood. Clinical manifestations of hypovolemic shock include hypotension and tachycardia. Therefore, the nurse should assess the client's status by obtaining her vital signs.

Scenario	Nurse Esther obtains Ms. Lieberman's vital signs after she reports feeling faint.
Question	Ms. Lieberman reports feeling worse and her vital signs are: BP 94/56 mm Hg, pulse 110/min, respirations 26/min, and SaO2 94%. Nurse Esther starts oxygen at 2 L/min. Which of the following should be Nurse Esther's priority action?
Selected Option	Recheck Ms. Lieberman's oxygen saturation.
Rationale	Ms. Lieberman's previous oxygen saturation level was below the expected reference range of 95% to 99%. After initiating oxygen therapy, the nurse should recheck the client's oxygen saturation. However this action is not the highest priority because it will not correct the underlying problem.

Optimal Decision	
Scenario	The unit of packed RBCs for Ms. Lieberman arrives on the unit.
Question	Nurse Esther is preparing to administer a unit of packed RBCs to Ms. Lieberman. Which of the following actions should Nurse Esther perform prior to administering the blood?

Selected Option	Ask Ms. Lieberman if she has experienced a reaction with any previous blood transfusions.
Rationale	A transfusion reaction can be caused by the development of antibodies to the donor leukocytes. This reaction is more likely to occur when a client has had blood transfusions before, as well as a history of prior blood transfusion reactions. Therefore, this is the appropriate action for the nurse to take.

Scenario	Nurse Esther is ready to administer the first unit of packed RBCs.
Question	Identify the correct sequence of actions for blood administration after Nurse Esther performs hand hygiene and applies gloves. (Reorder the steps by dragging them into the desired sequence.)
Selected Ordering	Gently rotate the bag of packed RBCs. Spike and prime the Y-set tubing with the 0.9% sodium chloride solution. Attach the packed RBCs bag to the Y-set tubing. Attach the tubing to the IV catheter and begin to infuse the 0.9% sodium chloride solution. Turn off the 0.9% sodium chloride solution. Begin to infuse the packed RBCs.
Rationale	The first action the nurse should do is insert one of the spikes of the Y-set into the 0.9% sodium chloride solution bag, prime the tubing with the 0.9% sodium chloride solution, and start slowly infusing the solution into the client's IV. Next, the nurse should gently rotate the bag to mix the blood cells with the plasma. Then, the nurse should spike the blood bag with the remaining spike on the Y-set tubing and turn off the 0.9% sodium chloride solution by closing the clamp. Lastly, the nurse needs to open the clamp to allow the blood to infuse.

Optimal Decision	
Scenario	Ms. Lieberman is restless, her face is flushed, and she reports having a headache.
Question	Nurse Esther notes Ms. Lieberman is restless, her face is flushed, and she reports having a headache. Her vital signs include: temperature 38.8° C (101.8° F), pulse 96/min, respirations 22/min, and BP 103/60 mm Hg. Which of the following is an appropriate action for Nurse Esther to take?
Selected Option	Stop the blood transfusion.
Rationale	In the presence of a febrile reaction, the client's blood is sensitive to some component of the donor's blood. To prevent further exposure to the sensitizing component, the transfusion should be stopped immediately.

Scenario	Using an SBAR format, write the information Nurse Esther should give to Ms. Lieberman's provider when calling about her response to the blood. Refer to the EMR documents for needed information.
Question	Using an SBAR format, write the information Nurse Esther should give to Ms. Lieberman's provider when calling about her response to the blood. Refer to the EMR documents for needed information. (Enter your response, then click on the submit button at the bottom of the screen. Compare your response to the one provided.)

Selected Option	S - Ms. Lieberman was admitted to our unit at 1500 for abd pain, blood stools, and dizziness. She was also admitted for transfusion of PRBCs and monitoring of stool. B - PRBC transfusion was started as soon as the blood was available on the unit. A - 15 mins into the PRBC transfusion, JL began to feel "chilly" and achy over her entire body. She is asking for Ibuprofen. Transfusion stopped as a result. T - 38.8 C, RR 22, BP 103/60, HR 96. R - Recommend prescribing antipyretic med.
Rationale	The following information should be shared with Ms. Lieberman's provider when calling about her response to the blood. Situation: Dr. McGuire, this is Esther - RN. I am taking care of Ms. Lieberman in room 5206. She is a 36-year-old client admitted from the ED today for a GI bleed. She's had one unit of packed RBCs and part of the second unit of blood. I stopped the second unit because I believe she is having a transfusion reaction. Her baseline temperature was 98.6 and is now 101.8. Ms. Lieberman reports having a headache, chills, and is restless. She does not have any evidence of a rash at this time. Background: Ms. Lieberman has a history of Crohn's disease and intermittent gastritis. Six months ago she had an ileostomy and started on infliximab IV every 8 weeks. Her last infusion was 7 weeks ago. Assessment: Her hemoglobin was 7 g/dL and her hematocrit was 21% in the ED. When she arrived to the medical surgical unit, her BP was 94/56 and her pulse 110, but now her BP is 110/70 and her pulse is 110. At this time, her respirations are 26, her SaO2 is 97%, and her temperature is 101.8. I have discontinued the second unit of blood and plan to send both the bags of blood to the lab per protocol. I hung a new bag of 0.9% sodium chloride to keep the line open. Recommendations: Ms. Lieberman is requesting ibuprofen for her headache, which would also bring her fever down. Could I have an prescription for an antipyretic, and do you want to continue the IV infusion of 0.9% sodium chloride at 150 mL/hr?

Scenario	Dr. March tells Ms. Lieberman that he recommends an endoscopy. Ms. Lieberman is informed about the procedure. She agrees to the procedure and signs the consent form.
Question	Nurse Esther is reinforcing teaching with Ms. Lieberman, who is scheduled for an endoscopy in the morning. Which of the following should Nurse Esther include in the teaching?
Selected Option	"A laxative will be given to clear the bowel."
Rationale	A laxative is unnecessary prior to an endoscopy. Additionally, a laxative is contraindicated for a client who has bleeding and cramping due to Crohn's disease.

Optimal Decision	
Scenario	Nurse Esther is calculating the number of milliliters of morphine sulfate to administer.
Question	Nurse Esther is preparing to administer morphine 4 mg IV. Available is morphine 8 mg/mL. How many mL should the nurse administer? (Round the answer to the nearest tenth.).
Selected Option	0.5

Rationale	Follow these steps for the Ratio and Proportion method of calculation: Step 1: What is the unit of measurement the nurse should calculate? mL Step 2: What is the dose the nurse should administer? Dose to administer = Desired 4 mg Step 3: What is the dose available? Dose available = Have 8 mg Step 4: Should the nurse convert the units of measurement? No Step 5: What is the quantity of the dose available? 1 mL Step 6: Set up an equation and solve for X. $\frac{\text{Have}}{\text{Desired}} = \frac{\text{Quantity}}{X} \Rightarrow \frac{8 \text{ mg}}{4 \text{ mg}} = \frac{1 \text{ mL}}{X \text{ mL}}$ $X \text{ mL} = 0.5 \text{ mL}$ Step 7: Round if necessary. Step 8: Determine whether the amount to administer makes sense. If there are 8 mg/mL and the prescription reads 4 mg, it makes sense to administer 0.5 mL. The nurse should administer morphine 0.5 mL IV. Follow these steps for the Desired Over Have method of calculation: Step 1: What is the unit of measurement the nurse should calculate? mL Step 2: What is the dose the nurse should administer? Dose to administer = Desired 4 mg Step 3: What is the dose available? Dose available = Have 8 mg Step 4: Should the nurse convert the units of measurement? No Step 5: What is the quantity of the dose available? 1 mL Step 6: Set up an equation and solve for X. $\frac{\text{Desired} \times \text{Quantity}}{\text{Have}} = \frac{4 \text{ mg} \times 1 \text{ mL}}{8 \text{ mg}} = \frac{0.5 \text{ mL}}{X \text{ mL}}$ $X \text{ mL} = 0.5 \text{ mL}$ Step 7: Round if necessary. Step 8: Determine whether the amount to administer makes sense. If there are 8 mg/mL and the prescription reads 4 mg, it makes sense to administer 0.5 mL. The nurse should administer morphine 0.5 mL IV. Follow these steps for the Dimensional Analysis method of calculation: Step 1: What is the unit of measurement the nurse should calculate? (Place the unit of measure being calculated on the left side of the equation.) $X \text{ mL} =$ Step 2: Determine the ratio that contains the same unit as the unit being calculated. (Place the ratio on the right side of the equation, ensuring that the unit in the numerator matches the unit being calculated.) $1 \text{ mL} \times \frac{8 \text{ mg}}{4 \text{ mg}} = 2 \text{ mL}$ Step 3: Place any remaining ratios that are relevant to the item on the right side of the equation, along with any needed conversion factors, to cancel out unwanted units of measurement. $1 \text{ mL} \times \frac{8 \text{ mg}}{4 \text{ mg}} = 2 \text{ mL}$ Step 4: Solve for X. $X \text{ mL} = 0.5 \text{ mL}$ Step 5: Round if necessary. Step 6: Determine whether the amount to administer makes sense. If there are 8 mg/mL and the prescription reads 4 mg, it makes sense to administer 0.5 mL. The nurse should administer morphine 0.5 mL IV.
------------------	---

Scenario	Nursing considerations Nurse Esther takes when administering morphine.
Question	Nurse Esther is preparing to administer 4 mg of morphine IV bolus to Ms. Lieberman. Which of the following actions should Nurse Esther take?
Selected Option	Inject the medication in the port on the IV tubing furthest from the IV insertion site.

Rationale	IV bolus medication should be injected in the port closest to the client.
------------------	---

Individual Report – Score Explanation and Interpretation

Reasoning Scenario Information:

Reasoning Scenario Information provides the date, time and duration of use, along with the score earned for each attempt. A Reasoning Scenario Performance score of Strong, Satisfactory, or Needs Improvement is provided for each attempt. This information is also provided for the Optimal Decision Mode if it has been enabled.

Reasoning Scenario Performance Scores:

Strong	Exhibits optimal reasoning that results in positive outcomes in the care of clients and resolution of problems.
Satisfactory	Exhibits reasoning that results in mildly helpful or neutral outcomes in the care of clients and resolution of problems.
Needs Improvement	Exhibits reasoning that results in harmful or detrimental outcomes in the care of clients and resolution of problems.

Reasoning Scenario Performance Related to Outcomes:

A clinical reasoning performance score related to each outcome is provided. Outcomes associated with student responses are listed in the report. The number across from each outcome indicates the percentage of responses associated with the level of performance of that outcome.

NCLEX® Client Need Categories:

Management of Care	Providing integrated, cost-effective care to clients by coordinating, supervising, and/or collaborating with members of the multi-disciplinary health care team.
Safety and Infection Control	Incorporating preventative safety measures in the provision of client care that provides for the health and well-being of clients, significant others, and members of the health care team.
Health Promotion and Maintenance	Providing and directing nursing care that encourages prevention and early detection of illness, as well as the promotion of health.
Psychosocial Integrity	Promoting mental, emotional, and social well-being of clients and significant others through the provision of nursing care.
Basic Care and Comfort	Promoting comfort while helping clients perform activities of daily living.
Pharmacological and Parenteral Therapies	Providing and directing administration of medication, including parenteral therapy.
Reduction of Risk Potential	Providing nursing care that decreases the risk of clients developing health-related complications.

Physiological Adaptation	Providing and directing nursing care for clients experiencing physical illness.
---------------------------------	---

Quality and Safety Education for Nurses (QSEN)

Safety	The minimization of risk factors that could cause injury or harm while promoting quality care and maintaining a secure environment for clients, self, and others.
Patient-Centered Care	The provision of caring and compassionate, culturally sensitive care that is based on a client's physiological, psychological, sociological, spiritual, and cultural needs, preferences, and values
Evidence Based Practice	The use of current knowledge from research and other credible sources, upon which clinical judgment and client care are based.
Informatics	The use of information technology as a communication and information gathering tool that supports clinical decision making and scientifically based nursing practice.
Quality Improvement	Care related and organizational processes that involve the development and implementation of a plan to improve health care services and better meet the needs of clients.
Teamwork and Collaboration	The delivery of client care in partnership with multidisciplinary members of the health care team, to achieve continuity of care and positive client outcomes.

Body Function

Cardiac Output and Tissue Perfusion	The anatomical structures (heart, blood vessels, and blood) and body functions that support adequate cardiac output and perfusion of body tissues.
Cognition and Sensation	The anatomical structures (brain, central and peripheral nervous systems, eyes and ears) and body functions that support perception, interpretation, and response to internal and external stimuli.
Excretion	The anatomical structures (kidney, ureters, and bladder) and body functions that support filtration and excretion of liquid wastes, regulate fluid and electrolyte and acid-base balance.
Immunity	The anatomic structures (spleen, thymus, bone marrow, and lymphatic system) and body functions related to inflammation, immunity, and cell growth.
Ingestion, Digestion, Absorption and Elimination	The anatomical structures (mouth, esophagus, stomach, gall bladder, liver, small and large bowel, and rectum) and body functions that support ingestion, digestion, and absorption of food and elimination of solid wastes from the body.
Integument	The anatomical structures (skin, hair, and nails) and body functions related to protecting the inner organs from the external environment and injury.
Mobility	The anatomical structures (bones, joints, and muscles) and body functions that support the body and provide its movement.

Oxygenation	The anatomical structures (nose, pharynx, larynx, trachea, and lungs) and body functions that support adequate oxygenation of tissues and removal of carbon dioxide.
Regulation and Metabolism	The anatomical structures (pituitary, thyroid, parathyroid, pancreas, and adrenal glands) and body functions that regulate the body's internal environment.
Reproduction	The anatomical structures (breasts, ovaries, fallopian tubes, uterus, vagina, vulva, testicles, prostate, scrotum, and penis) and body functions that support reproductive functions.

Decision Log

Information related to each question answered in a scenario attempt is listed in the report. A brief description of the scenario, question, selected option and rationale for that option are provided for each question answered. The words "Optimal Decision" appear next to the question when the most optimal option was selected.

The rationale for each selected option may be used to guide remediation. A variety of learning resources may be used in the review process, including related ATI Review Modules.