

<p style="text-align: center;">Anticipated Patient Problem</p> <p style="text-align: center;">and</p> <p style="text-align: center;">Goals</p>	<p style="text-align: center;">Relevant Assessments</p> <p style="text-align: center;">(Prewrite) What assessments pertain to your patient's problem? Include frequencies</p>	<p style="text-align: center;">Multidisciplinary Team Intervention</p> <p style="text-align: center;">(Prewrite) What will you do if your assessment is abnormal?</p>
<p>Problem: Impaired Urinary Elimination</p> <p>Reasoning: decreased urine output, acute kidney injury</p> <p>Goal: Patient will void 30ml/hr during my time of care.</p> <p>Goal: Patient will demonstrate proper bladder emptying techniques, to prevent urinary retention and reduce the risk of urinary tract infections evidenced by maintaining a post-void residual volume of less than 50ml and reporting clear, odor free urine during my time of care.</p>	<p>Assess urine characteristics q4hrs or PRN</p>	<p>Maintain patency of indwelling catheter PRN during my time of care.</p>
	<p>Assess urine output q4hr or PRN</p>	<p>Administer diuretics per order during my time of care.</p>
	<p>Assess urine after each void or PRN</p>	<p>Educate client on importance of straining urine and how to strain urine during my time of care.</p>
	<p>Assess patients voiding schedule during my time of care.</p>	<p>Encourage toileting q2hr to create a schedule during my time of care.</p>
	<p>Monitor BUN, creatinine, and WBC daily</p>	<p>Maintain renal function-encourage UA dipstick as needed</p>
	<p>Assess willingness to learn new techniques to strengthen muscles during my time of care.</p>	<p>Educate and encourage the use of Kegel exercises q2hr or PRN during my time of care.</p>

Additional problems: Risk for unstable blood glucose, Impaired gas exchange

<p style="text-align: center;">Anticipated Patient Problem</p> <p style="text-align: center;">and</p> <p style="text-align: center;">Goals</p>	<p style="text-align: center;">Relevant Assessments</p> <p style="text-align: center;">(Prewrite) What assessments pertain to your patient's problem? Include frequencies</p>	<p style="text-align: center;">Multidisciplinary Team Intervention</p> <p style="text-align: center;">(Prewrite) What will you do if your assessment is abnormal?</p>
<p>Problem: Risk for bleeding</p> <p>Reasoning: hematocrit 36.8, hemoglobin 12.4, three blood products given in ICU, PTT 40, PT time 13</p> <p>Goal: Patient will not experience bleeding as evidence by normal blood pressure of 120/80, an increasing hemoglobin level of 13.2 and a hematocrit of at least 40 during my time of care.</p> <p>Goal: Patient will demonstrate measures to prevent bleeding such as using electric razor, soft toothbrush, no flossing and recognize signs of bleeding that need to be reported immediately to the provider such as dark tarry stool during my time of care.</p>	<p>Assess Hemoglobin and Hematocrit levels daily</p>	<p>Administer blood products as needed with decreasing H&H levels during my time of care.</p>
	<p>Assess PT time and PTT daily</p>	<p>Administer Heparin 5,000 units q8hr during my time of care</p>
	<p>Monitor vital signs- BP and HR q4hr or PRN</p>	<p>Maintain stable BP and HR Educate patient on need for IV bolus if BP drops due to bleeding during my time of care.</p>
	<p>Assess Stool and urine PRN for bleeding</p>	<p>Educate patient on how to check stool for bleeding, dark tarry stool.</p>
	<p>Assess skin for signs of petechiae, bruising, hematoma formation daily</p>	<p>Educate patient on decreasing bleeding risk and when to report, such as petechiae, bruising.</p>
	<p>Assess client's medication history (anticoagulants) and health history increasing risk of bleeding</p>	<p>Educate client on the use of medications that can increase risk for bleeding such as anticoagulants during my time of care.</p>