

ALEXA DOIAN

Immunology Class Preparation

- Label each example as either active or passive immunity:
 - Antibodies are passed from the mother to the fetus: passive
 - Antibodies are produced after exposure to a killed virus: Active
 - Antibodies are produced after an infection: Active
 - Antibodies are administered in the form of immune globulins: passive
- A nurse is assessing a client for HIV. The nurse should identify which of the following risk factors associated with this virus? Select all that apply.
 - Perinatal exposure
 - Monogamous partner
 - Blood transfusion
 - Occupational exposure
- A nurse in an outpatient clinic is assessing a clinic who reports night sweats, fatigue, cough, nausea, diarrhea and has a temperature of 38.1° C (100.6° F). The client is concerned about the possibility of having HIV. Which actions should the nurse take? Select all that apply.
 - Perform a physical assessment
 - Determine when the manifestations began
 - Request a prescription for an antibiotic
 - Request a prescription for HIV testing
 - Obtain a sexual history
- List three effects of aging on the immune system:
 - Ability to fight off infection/cancer decreases.
 - increased susceptibility of autoimmune disorders
 - Reduced effectiveness of vaccines.
- Match the type of hypersensitivity reaction to the characteristics:

Type I: IgE-Mediated <u>A</u>	<u>A.</u> Caused by pollen, food, drugs, dust, immediate reaction, allergic rhinitis, atopic dermatitis, skin test shows wheal and flare
Type II: Cytotoxic <u>C</u>	<u>B.</u> Occurs when T cells attack antigens or release cytokines, several days to occur, contact dermatitis
Type III: Immune-Complex <u>D</u>	<u>C.</u> Caused by ABO incompatibility blood transfusion reaction, occurs in minutes to hours, activates
Type IV: Delayed Hypersensitivity <u>B</u>	<u>D.</u> Caused by fungal, viral, or bacterial antigens, takes hours to days to develop, rheumatoid arthritis, systemic lupus erythema complement system and leads to cell lysis and acute kidney injury