

Cardiac Assessment Definition Match

- | | |
|---------------------------------------|--|
| H. Thrill | F. Turbulent sounds occurring between normal heart sounds. Characterized by loudness, pitch, shape, duration. Graded I-VI |
| K. 3 rd Heart Sound | D. Small red streaks under fingernails |
| I. Troponins | G. HR > 100 |
| N. HDL | E. HR < 60 |
| F. Heart murmurs | B. The amount of blood pumped by each ventricle in 1 min. Calculated by Stroke Volume X HR |
| D. Splinter hemorrhages | M. Peripheral resistance against which the left ventricle must pump |
| G Tachycardia | A. Turbulent flow through artery caused by aneurysm or arterial obstruction |
| E. Bradycardia | I. Contractile proteins released after MI. Both T and I samples test for cardiac injury/ischemia |
| A Arterial bruit | K. Extra heart sound heard in EARLY diastole, similar to the sound of a gallop. Causes include Left Ventricular failure, volume overload, aortic or tricuspid regurgitation |
| J LDL | L. Alteration of spine often in elderly resulting in altered chest landmarks, auscultation, and percussion |
| B Cardiac Output | C. The volume of blood stretching the ventricles at the end of diastole before the next contraction |
| L Kyphosis | H. Vibration of vessel or chest |
| C Preload | N. Lipoprotein considered to be the “good cholesterol” |
| M Afterload | J. Lipoprotein considered to be the “lethal cholesterol” |