

Actual Patient Problem: hemorrhage

Clinical Reasoning: placental wound, vaginal delivery, 1st degree laceration, full bladder

Goal: Fundal height will be appropriate (-1 on day 1) postpartum, firm, and midline

Goal: Lochia will be rubra on day 1-3 with <500mL CBL during my TOC

Met: Unmet:

Met: Unmet:

Actual Patient Problem: R/F infection

Clinical Reasoning: vaginal delivery, midline laceration, homeless

Goal: will not have uterine s/s of infection as evidence by appropriate fundal height, location, and tone for days pp, appropriate amount of lochia, and no foul smelling lochia during my time of care

Met: Unmet:

Goal: will not have any s/s of systemic signs of infection and will have a temp between 36.1-37.2 degrees, and a HR between 60-100bpm during my time of care

Met: Unmet:

Patient Problem	Time	Relevant Assessments Indicate pertinent assessment findings.	Time	Multidisciplinary Team Intervention What interventions were done in response to your abnormal assessments?	Time	Reassessment/Evaluation What was your patient's response to the intervention?
Bleeding/risk for	1000	Lochia moderate/rubra, BP: 115/75, HR: 96, fundus firm/midline/at umbilicus saturated chuck pad, midline laceration/well-approximated	1010	Weighed pad, placed a clean and dry pad down	1020	New chuck pad in place, continuing to bleed moderately, blood loss: 118mL
Bleeding	1010	Boggy uterus w/ +2, lochia rubra, complains of feeling trickling in the pad, blood loss: 118mL	1015	Started fundal massage	1025	fundus boggy at midline +1, lochia continues trickling, blood loss: 118mL
Bleeding	1010	Total blood loss: 518mL, HR: 110, continuing to bleed/saturated pad, BP: 99/50, HR: 120	1015	Called the provider about increasing HR, total blood loss of 518mL with continued bleeding	1020	Continued to massage the fundus, postpartum hemorrhage orders placed, called the provider, BP: 95/50, HR: 120
Bleeding	1020	Provider placed post-partum hemorrhage orders, BP: 95/50, HR:121	1030	Administered oxytocin IV at 500ml/hr (30 units/hr), 1000ml IV bolus of NS fluid, 0.2mg of methergine IM	1050	Continued to monitor blood loss, total blood loss: 955mL, continued to massage the fundus
Bleeding	1025	Oxygen 92% on RA, client complains of dizziness and malaise	1025	Flattened HOB, applied nonrebreather at 100% oxygen	1120	Oxygen 98% with non-rebreather on, stated dizziness alleviating

Risk for infection	1045	Fundus boggy at midline +1, displaced to the right, has not been able to use the bathroom since this morning	1050	Placed straight cath using a sterile technique	1100	4
Hemorrhage	1120	fundus firm and midline after emptying bladder, BP increased to 98/58, no longer complains of dizziness	1125	Changed pads and sheet, raised the HOB, weighed the pad	1130	1

Significant Event Documentation: Use the area below to document any significant events that happened during your time of care.
