

R.B. Turnbull, Jr. MD School of WOC Nursing Education

Ostomy Care Mini Case Studies



Student Name & Date: _____

Reviewed by _____

Score: /46

This assignment focuses on applying the assessment of an individual with an ostomy to pouching principles. First, basic principles are identified. Then, principles are applied to clinical situations.

Answer the following questions:

1. Identify the nursing orders for changing a pouching system on a person with no peristomal skin breakdown. (2 points)
2. Identify nursing orders for changing a pouching system on a person with peristomal skin breakdown. (2 points)
3. Identify nursing orders for changing a pouching system on a person with peristomal skin breakdown and the presence of satellite lesions. (2 points)

For each of the below ostomy patient case scenarios:

- ❖ Use the information provided to identify an ostomy pouching plan.
 - ❖ Be specific: It is important to note a pouching system is a skin barrier wafer and a pouch. A complete answer should include both unless otherwise indicated. **Include the manufacturer and full product name.** Product numbers should not be used. Make sure to include accessory products as needed.
 - ❖ When providing the rationale: Describe abdominal characteristics, stoma characteristics, and one other reason why you would choose the specific system.
- ❖ The first half of the first case study has been completed for you below as an example:

Example + Scenario 1



55-year-old with a history of colon cancer. Colostomy was created 2 months ago and presents today in the ostomy clinic for assessment and management. Pt is very active and would like to consider a more flexible pouching system. Pt is changing his pouching system every other day because he is fearful of leakage.

Assessment: Stoma is pink, budded, and protrudes above skin level. No erythema on peristomal skin. No reports of leakage.

Identify a one and two-piece pouching system option along with rationale for choice.

Image courtesy of Wound, Ostomy, and Continence Nurses Society™ image library.

One Piece System: *Hollister Premier one-piece drainable pouch flat Flexwear barrier with clamp closure, change every 5-7 days and PRN.*

Rationale: *This system is flexible and matches the contours of this patient's abdomen. It is appropriate for budded stomas with an even peristomal plane and is manufactured for wear for multiple days.*

Two Piece option:

Rationale:

/2 points

Scenario 2



42-year-old with stoma placement on soft, obese abdomen.

Assessment: Stoma pink, budded, and protruding. Edema and necrosis circumferential at stomal edge. Serosanguineous drainage in pouch. Skin barrier wafer removal notes being cut too small, restricting and causing trauma to the stoma.

Identify a one and two-piece pouching system option along with rationale for choice.

Image courtesy of Wound, Ostomy, and Continence Nurses Society™ image library.

One Piece option:

Rationale:

Two Piece option

Rationale:

/4 points

Scenario 3



85-year-old presents with flush ileostomy and peristomal irritant dermatitis. Oval stoma with os at 6 o'clock location. Protuberant hernia above further pushes the stoma into a lateral fold.

Pt wears bifocal glasses when applying the pouching system. Due to extreme hip contours, it is difficult to have a hernia belt stay in place.

Image courtesy of Wound, Ostomy, and Continence Nurses Society™ image library.

Pouching recommendations:

Rationale:

/2 points

Scenario 4



56-year-old obese individual with ruptured diverticulitis. A red rubber catheter in place as a bridge for the loop ostomy. Stoma is slightly budded and red. Peristomal skin with erythema and partial thickness wound 4-7 o'clock Etiology may be due to trauma from red rubber catheter movement.

Image courtesy of Wound, Ostomy, and Continence Nurses Society™ image library.

Pouching recommendations:

Rationale:

/2 points

Scenario 5



42-year-old arrives in emergency room with complaints of difficulty pouching and peristomal skin irritation. Current pouching system sometimes has less than 4 hours of wear time. Skin is very painful. Assessment finding of ulcerated skin around stoma. Stoma is at skin level on a firm abdomen. Patient acknowledges frequent sweating resulting in the need to change appliance. "It just doesn't seem to stick".

Image courtesy of Wound, Ostomy, and Continence Nurses Society™ image library.

Pouching Recommendations:

Rationale:

/2 points

Scenario 6



66-year-old obese individual with stoma in an abdominal fold. Appliance leakage causing contact dermatitis. Wear time has been less than 8 hours. Irritation is painful.

Image courtesy of Wound, Ostomy, and Continence Nurses Society™ image library.

Pouching Recommendations:

Rationale:

/2 points

Scenario 7



76-year-old presents to the ostomy clinic with peristomal redness to periphery. Irritation limited to appliance tape collar region. Satellite lesions present. Stoma is budded and round. States has had ostomy for 6 months and has not had any problem until recently after Home Health changed the products.

Image courtesy of Wound, Ostomy, and Continence Nurses Society™ image library.

Pouching Recommendations:

Rationale:

/2 points

Scenario 8



Individual presents to the clinic with stoma measuring 3.5 inches. Stoma protrudes above skin level. Uneven peristomal contours with skin folds at 3 and 9 o'clock. Moisture-related skin damage on peristomal skin related to leakage.

Image courtesy of Wound, Ostomy, and Continence Nurses Society™ image library.

Pouching Recommendations:

Rationale:

/2 points

Scenario 9



Patient presents to ostomy clinic due to peristomal hernia causing peristomal skin breakdown. Abdomen is firm. Appliance wear time has decreased since parastomal hernia development. Stoma is flush with skin. Os between 5 and 6 o'clock area. Complains of odor. "The odor is really bad when I empty the pouch".

Image courtesy of Wound, Ostomy, and Continence Nurses Society™ image library.

Pouching Recommendations:

Rationale:

Odor Management Strategies:

/3 points

Scenario 10



A pediatric Individual presents to the emergency room with stoma prolapse. Caregiver expresses inability to apply pouching system related to stomal protrusion. Stoma is red and healthy. No peristomal irritation.

Identify one pouching system with rationale for choice along with one consideration with appliance application specific to a prolapsed stoma.

Image courtesy of Wound, Ostomy, and Continence Nurses Society™ image library.

Pouching Recommendations:

Rationale:

Further Considerations:

/3 points

Scenario 11



A 28-year-old with an ileostomy presents to the clinic for a follow-up evaluation. During the visit, the patient expressed the pouch is too long with the end of the pouch falling into the groin area. Assessment notes stoma red, viable, and protrudes above skin level. Abdominal space is small with short distance from stoma to groin. Current appliance is a one-piece cut to fit skin barrier. Pouch length 12". Name at least two alternative pouching management system options and rationale for each.

Image courtesy of Judy Mosier, MSN, RN, CWOCN

Pouching option #1:

Rationale:

Pouching option #2:

Rationale:

/4 points

Scenario 12



You are in your office and take a call from a patient. The patient voices having to change the skin barrier wafer more frequently, itching under the skin barrier, and desire to change manufacturers. The patient agrees to be seen in the clinic.

In preparation for this visit, you go to your resources to help you.

1. Identify one manufacturer (Hollister, Convatec, Coloplast, NuHope, etc)
2. Identify three skin barrier wafers from that manufacturer that differ in composition/ingredients.
3. Identify the type of ostomy or situation in which the wafer is appropriate.

For example: (can not be used)

Manufacturer: B. Braun

1. Skin barrier wafer: Flexima 3S

Composition & Purpose: Made of new generation plastics making it more soft and flexible. Appropriate for any type of ostomy and active individuals

2. Skin barrier wafer: Flexima... etc

Manufacturer:

Skin barrier Wafer 1:

Composition & Purpose:

Skin barrier Wafer 2:

Composition & Purpose:

Skin barrier Wafer 3:

Composition & Purpose:

/6 points

Scenario 13




Image courtesy of Judy Mosier, MSN, RN, CWOCN

You are consulted to see a patient with a new colostomy. Upon entering the room, you note there is an indwelling catheter in the stoma. Nursing reports pouch leakage as the hole in the pouch for the tube is cut to fit the stoma resulting in a “big hole” in the front of the pouch. The surgeon’s request is to continue to pouch the stoma while pulling the tube through the pouch.

Describe how you will secure the tube while separately pouching the stoma and the tube

...using a commercial access port:

...in the absence of a commercial access port:

/2 points

Scenario 14



86-year-old obese individual presents to the ostomy clinic with a retracted stoma. States has a soft-formed stool once a day. Pouch changed daily as stool goes under the skin barrier wafer, and at times, no stool goes into the pouch.

It is determined a convex pouching system should be used. A convex skin barrier wafer is not available.

Identify two strategies to create convexity in the absence of a convex skin barrier wafer.

Image courtesy of Wound, Ostomy, and Continence Nurses Society™ image library.

Alternative convexity option #1:

Alternative convexity option #2:

/2 points

Scenario 15



The WOC nurse is consulted to manage a wound with a stoma in proximity. The surgeon has consented to pouching the stoma in the same pouch as the wound. It is determined to be the best approach.


Identify one product that can be used to achieve this.

Image courtesy of Judy Mosier, MSN, RN, CWOCN

Pouching option:

/1 point

Scenario 16



A 70-year-old patient presents to the ED with pouching difficulty. They report using a fistula pouch previously, however, this has become too costly of an option. Their stoma measures 4 1/3" in diameter and they are at a loss for pouching options. The patient will need pouching long term. Identify one product pouching system that is manufactured to accommodate a stoma of 4" or greater in size.

Image courtesy of Dr. James Wu

Pouching option:

/1 point