

Braden Scale for Predicting Pressure Sore Risk

Instructions:

Use the Braden Scale to assess the patient's level of risk for development of pressure ulcers. The evaluation is based on six indicators: sensory perception, moisture, activity, mobility, nutrition, and friction or shear.

Scoring:

The Braden Scale is a summated rating scale made up of six subscales scored from 1-3 or 4, for total scores that range from 6-23. A lower Braden Scale Score indicates a lower level of functioning and, therefore, a higher level of risk for pressure ulcer development. A score of 19 or higher, for instance, would indicate that the patient is at low risk, with no need for treatment at this time. The assessment can also be used to evaluate the course of a particular treatment.

Sources:

- Ayello EA. Predicting pressure ulcer risk. In: Boltz M, series ed. *Try This: Best Practices in Nursing Care to Older Adults*. 2003 July. Revised January 2004; Vol 1, No 5. The Hartford Institute for Geriatric Nursing. www.hartfordign.org
- Ayello EA, Braden B. How and why to do pressure ulcer risk assessment. *Adv Skin Wound Care*. 2002 May-Jun;15(3):125-131.
- Bergstrom N, Braden BJ, Laguzza A, Holman V. The Braden Scale for Predicting Pressure Sore Risk. *Nurs Res*. 1987;36:205-210.
- Braden Scale for Preventing Pressure Sore Risk. Prevention Plus. 2001. Available at: <http://www.bradenscale.com/bradenscale.htm>. Accessed December 16, 2004.
- Panel for the Prediction and Prevention of Pressure Ulcers in Adults. Pressure Ulcers in Adults: Prediction and Prevention. Clinical Practice Guideline, Number 3. AHCPR Publication No. 92-0047. Rockville, MD: Agency for Health Care Policy and Research, Public Health Service, U.S. Department of Health and Human Services. May 1992 (reviewed 2000).

BRADEN SCALE FOR PREDICTING PRESSURE SORE RISK

Last two digits of Patient's room number _____ Evaluator's Name _____

Date of Assessment

SENSORY PERCEPTION ability to respond meaningfully to pressure-related discomfort	1. Completely Limited Unresponsive (does not moan, flinch, or grasp) to painful stimuli, due to diminished level of consciousness or sedation OR limited ability to feel pain over most of body.	2. Very Limited Responds only to painful stimuli. Cannot communicate discomfort except by moaning or restlessness OR has a sensory impairment which limits the ability to feel pain or discomfort over ½ of body.	3. Slightly Limited Responds to verbal commands, but cannot always communicate discomfort or the need to be turned OR has some sensory impairment which limits ability to feel pain or discomfort in 1 or 2 extremities.	4. No Impairment Responds to verbal commands. Has no sensory deficit which would limit ability to feel or voice pain or discomfort.				
MOISTURE degree to which skin is exposed to moisture	1. Constantly Moist Skin is kept moist almost constantly by perspiration, urine, etc. Dampness is detected every time patient is moved or turned.	2. Very Moist Skin is often, but not always moist. Linen must be changed at least once a shift.	3. Occasionally Moist Skin is occasionally moist, requiring an extra linen change approximately once a day.	4. Rarely Moist Skin is usually dry, linen only requires changing at routine intervals.				
ACTIVITY degree of physical activity	1. Bedfast Confined to bed.	2. Chairfast Ability to walk severely limited or non-existent. Cannot bear own weight and/or must be assisted into chair or wheelchair.	3. Walks Occasionally Walks occasionally during day, but for very short distances, with or without assistance. Spends majority of each shift in bed or chair.	4. Walks Frequently Walks outside room at least twice a day and inside room at least once every two hours during waking hours.				
MOBILITY ability to change and control body position	1. Completely Immobile Does not make even slight changes in body or extremity position without assistance.	2. Very Limited Makes occasional slight changes in body or extremity position but unable to make frequent or significant changes independently.	3. Slightly Limited Makes frequent though slight changes in body or extremity position independently.	4. No Limitation Makes major and frequent changes in position without assistance.				
NUTRITION usual food intake pattern	1. Very Poor Never eats a complete meal. Rarely eats more than ½ of any food offered. Eats 2 servings or less of protein (meat or dairy products) per day. Takes fluids poorly. Does not take a liquid dietary supplement OR is NPO and/or maintained on clear liquids or IVs for more than 5 days.	2. Probably Inadequate Rarely eats a complete meal and generally eats only about ½ of any food offered. Protein intake includes only 3 servings of meat or dairy products per day. Occasionally will take a dietary supplement OR receives less than optimum amount of liquid diet or tube feeding.	3. Adequate Eats over half of most meals. Eats a total of 4 servings of protein (meat, dairy products) per day. Occasionally will refuse a meal, but will usually take a supplement when offered OR is on a tube feeding or TPN regimen which probably meets most of nutritional needs.	4. Excellent Eats most of every meal. Never refuses a meal. Usually eats a total of 4 or more servings of meat and dairy products. Occasionally eats between meals. Does not require supplementation.				
FRICITION & SHEAR	1. Problem Requires moderate to maximum assistance in moving. Complete lifting without sliding against sheets is impossible. Frequently slides down in bed or chair, requiring frequent repositioning with maximum assistance. Spasticity, contractures or agitation leads to almost constant friction.	2. Potential Problem Moves feebly or requires minimum assistance. During a move skin probably slides to some extent against sheets, chair, restraints or other devices. Maintains relatively good position in chair or bed most of the time but occasionally slides down.	3. No Apparent Problem Moves in bed and in chair independently and has sufficient muscle strength to lift up completely during move. Maintains good position in bed or chair.					
					Total Score			