

## Reconstitution for IVP

Medication	Reconstitution	Diluent	Comments	Rate of IV Push Administration
<b>Acetazolamide (Diamox)</b>	500 mg in 5 ml	Sterile Water	Shake vigorously	3-5 minutes
<b>Alteplase (Cathflo)</b>	2 mg in 2 ml	2.2 ml Sterile Water	<b>DO NOT SHAKE</b> (gently swirl)	over 1 minute
<b>Aztreonam (Azactam)</b>	1 g in 10 ml 2 g in 10 ml	Sterile Water	Shake vigorously	3-5 minutes
<b>Calcium Gluconate</b>	1 g/10 ml (liquid)	Undiluted	N/A	Over 10 minutes
<b>Cefazolin (Ancef)</b>	1 g in 10 ml 2 g in 20 ml	Sterile Water	Shake vigorously	3-5 minutes
<b>Cefotetan (Cefotan)</b>	1 g in 10 ml 2 g in 20 ml	Sterile Water	Shake vigorously	3-5 minutes
<b>Cefuroxime (Zinacef)</b>	750 mg in 10 ml 1.5 g in 20 ml	Sterile Water or NS	Shake vigorously	3-5 minutes
<b>Ceftriaxone (Rocephin)</b>	1 g in 10 ml 2 g in 20 ml	Normal Saline	Shake vigorously	3-5 minutes
<b>Ceftazidime (Tazicef, Fortaz)</b>	1 g in 10 ml 2 g in 20 ml	Sterile Water	Shake vigorously	3-5 minutes
<b>Cefepime (Maxipime)</b>	1 g in 10 ml 2 g in 20 ml	Sterile Water or NS	Shake vigorously	3-5 minutes
<b>Methocarbamol (Robaxin)</b>	1 g/10 ml (liquid)	Undiluted	N/A	3-5 minutes
<b>Meropenem (Merrem)</b>	500 mg in 10 ml 1 g in 20 ml	Sterile Water	Shake vigorously	3-5 minutes
<b>Pantoprazole (Protonix)</b>	40 mg in 10 ml	Normal Saline	Shake vigorously	2-4 minutes

**IVP medications are to be given immediately, not to exceed 1 hour after reconstitution.**

Not all-inclusive list, call pharmacy for questions.