

IM7 Lab Week 2

Titration Lab Quiz

1. Which of the following are true about norepinephrine (Levophed)? Select all that apply
 - a. Norepinephrine may cause dysrhythmias
 - b. Extravasation may cause tissue necrosis
 - c. Norepinephrine dilates the veins and arteries
 - d. Norepinephrine may increase O₂ demand

2. A patient is receiving a continuous propofol infusion. Which of the following should the nurse monitor as a potential adverse effect? Select all that apply
 - a. Respiratory depression
 - b. Hypertension
 - c. Elevated triglyceride level
 - d. Acidosis
 - e. Rhabdomyolysis

3. While assessing a patient receiving nicardipine infusion the nurse should expect to find which of the following if the infusion is effective?
 - a. Decreased blood pressure
 - b. Decreased heart rate
 - c. Increased serum calcium level
 - d. Normal respiratory rate

Chest Tube Quiz - Review the assigned HESI Skills on Chest Tube:

4. You are providing care to a patient with a chest tube. On assessment of the drainage system, you note continuous bubbling in the water seal chamber and oscillation. Which of the following is the correct nursing intervention for this type of finding?
 - a. Reposition the patient because the tubing is kinked.
 - b. Continue to monitor the drainage system.
 - c. Increase the suction to the drainage system until the bubbling stops.
 - d. Check the drainage system for an air leak.

5. A patient calls and tells you something is wrong with his chest tube. When you arrive to the room you note that the drainage system has fallen on its side and is leaking drainage onto the floor from a crack in the system. What should be your next action?
 - a. Place the patient in supine position and clamp the tubing
 - b. Notify the physician immediately
 - c. Disconnect the drainage system and get a new one
 - d. Disconnect the tubing from the drainage system and insert the tubing 1 inch into a bottle of sterile water and obtain a new system

12. Once a code is called and CPR initiated, what things justify interrupting compressions? Select all that apply

- i. Rhythm or pulse check
- ii. Shock
- iii. Ventilations
- iv. Medication administration

13. What joule setting does the manufacturer recommend as the start for the Biphasic defibrillator?

- i. 50-100
- ii. 120-200
- iii. 200-300
- iv. 300-360