

**Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Oxytocic	Induce labor, postpartum hemorrhage	IM IV	Y N		Maternal- N/V, COMA, SEIZURES, hypotension  Fetal- INTRACRANIAL HEMORRHAGE, hypoxia	1. Fetal maturity, presentation, and pelvic adequacy should be assessed prior to administration of oxytocin for induction of labor 2. Monitor maternal BP and pulse frequency and fetal heart rate continuously throughout administration 3. Monitor maternal electrolytes 4. Monitor contractions
Magnesium Sulfate	Mineral/ electrolyte	Mineral & electrolyte replacements/ supplements	IV PO IM	Y N		Drowsiness, decreased respiratory rate, flushing, sweating, muscle weakness	1. Monitor for toxemia 2. Monitor newborn for hypotension and respiratory depression 3. Monitor input and output 4. Take serum magnesium as ordered
Meperidine	Opioid agonists	Pain	IM IV PO	Y N		Confusion, sedation, hypotension, constipation, N/ V	1. Monitor pain level 2. Monitor for respiratory distress 3. Teach patient about possible constipation 4. Teach patient to change positions to avoid orthostatic hypotension
Promethazine	Tricyclic	Relaxation during labor	IM IV	Y N		Drowsiness, dizziness, constipation,	1. Monitor respiratory status and oxygen levels 2. Teach patient to avoid prolonged exposure to sunlight

Student Name: \_\_\_\_\_

Unit: \_\_\_\_\_

Pt. Initials: \_\_\_\_\_

Date: \_\_\_\_\_

**Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

						blurred vision, dry mouth	3. Can cause drowsiness- call before getting out of bed 4. Ensure patient has non-slip socks to prevent falls
Calcium Gluconate	Calcium Supplement	Increase calcium levels	IV PO	Y N		Edema, lightheadedness, bradycardia, hypercalcemia, confusion	1. Monitor serum Calcium levels 2. Rapid IV infusion can cause hypotension, bradycardia, or cardiac arrest 3. Monitor heart rate and BP closely 4. Monitor fluid and electrolytes closely to avoid fluid overload

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	Vitamin	Prevent bleeding	IV Oral Sub Q	Y N		Taste changes, tachycardia, SOB, cyanosis, flushed skin	1. Monitor electrolyte levels 2. Monitor respiratory status closely 3. Monitor signs/symptoms of bleeding 4. Teach the patient to inform the nurse of any abnormal taste they may be experiencing
Erythromycin Ophthalmic Ointment	Antibiotic	Prophylaxis for bacterial infection of eyes	Topical	Y N		Eye irritation, hypersensitivity, redness, vision changes	1. Apply to newborn's eyes within first hour after birth 2. Instill 1 cm across newborns eyes 3. Do not touch tip of applicator to eye 4. This ointment may increase the serum concentration of Vitamin K antagonists
Engerix B	Vaccine	Hepatitis B;	10mcg/ 0.5 mL	Y		Injection site	1. Shake well before withdrawal and use

**Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

		Prophylaxis	IM Sub Q	N		reaction, HA, Malaise, Nausea, Diarrhea, Anaphylaxis, Pancytopenia	2. Administer in the deltoid area for older children and anterolateral thigh in infants and young children 3. Engerix- B is given in a series of 2-4 injections 4. Teach patient about dosing schedule
Hepatitis B Immune Globulin	Immune Serum	Hepatitis B; Prophylaxis	0.5 mL IM IV	Y N		Erythema at injection site, injection site pain, N/V, decreased WBC	1. Give to newborn within 12 hours of birth 2. Avoid live vaccines other than Hep B vaccine for 3 months after receiving this drug 3. Drug may cause thrombotic events 4. This drug may interfere with some blood glucose monitoring systems, and result in falsely elevated glucose reading
				Y N			Sources: Micromedex App Multum, C. (2020, August 19). Erythromycin ophthalmic Uses, Side Effects & Warnings. Retrieved October 06, 2020, from <a href="https://www.drugs.com/mtm/erythromycin-ophthalmic.html">https://www.drugs.com/mtm/erythromycin-ophthalmic.html</a>