

## Dosage Calculation Worksheet #2

1. The IV order is for D<sub>5</sub>W to infuse at 100 mL/hr. The drop factor is 10 gtt/mL. How many drops per minute (gtt/min) should the pump be set to run? Round final answer to whole number.

$$\frac{V}{T} \times \text{gtt/mL} = Y$$

$$\frac{100}{60} = 1.6666 \times 10 \text{ gtt/mL} = 16.66 = 17 \text{ gtt/min}$$

2. Medication order: Rocephin 1 g IV every 12 hours over 30 minutes. Available: rocephin 1 g in 150 mL NS. At what rate would you set your pump?

$$\frac{V}{T} = Y$$

$$\frac{30}{60} = 0.5 \quad \frac{150}{0.5} = 300 \text{ mL/hr}$$

3. Medication order: Vistaril 20 mg IM q4h PRN for nausea. The 10 mL vial that you have available is labeled 25 mg/mL. How many mL will you draw up to give?

$$\frac{25 \text{ mg}}{1 \text{ mL}} = \frac{20 \text{ mg}}{X} = 0.8 \text{ mL}$$

4. Medication order: Haldol 3 mg IM q6h PRN for agitation. The 1 mL vial that you have available is labeled 5 mg/mL. How many mL will you draw up to give?

$$\frac{5 \text{ mg}}{1 \text{ mL}} = \frac{3 \text{ mg}}{X} = 0.6 \text{ mL}$$

5. Medication order: heparin 5000 units subQ every 12 hours. Drug available: heparin 10,000 units/2 mL. How many mL will you administer for the day?

$$\frac{10000}{2 \text{ mL}} = 5000 \text{ units/mL} \cdot \frac{5000}{5000} = 10 \text{ mL}$$

6. A patient has an order for 200 mg q8h of cimetidine (Tagamet) to be administered intramuscularly. The vial of 8 mL contains 300 mg per 2 mL. How many mL would you give q8h?

$$\frac{300 \text{ mg}}{2 \text{ mL}} = \frac{200 \text{ mg}}{X} = 1.3 \text{ mL}$$

7. Medication order: Garamycin 80 mg IVPB over 30 minutes. Available: Garamycin (gentamicin sulfate) 80 mg in 50 mL of D<sub>5</sub>W. Calculate the flow rate in mL/hr.

$$\frac{30}{60} = 0.5 \quad \frac{50}{0.5} = 100 \text{ mL/hr}$$

8. You have an IV infusing at 125 mL/hr. How long will it take 1500 mL to infuse?

$$\frac{V}{T} = r \quad \frac{1500}{125} = 12 \text{ hrs}$$

Same as #2?

9. Medication order: rocephin 1 g IV every 12 hours over 30 minutes. Available: rocephin 1 g in 150 mL NS. At what rate would you set your pump?

$$\frac{30}{60} = 0.5 \quad \frac{150}{0.5} = 300 \text{ mL/hr}$$

10. An infusion pump is set to administer 75 mL/hr to a patient. How many hours will it take for the patient to receive 600 mL of fluid?

$$\frac{600}{75} = 8 \text{ hrs}$$

11. A patient is to receive lidocaine hydrochloride (Xylocaine) 100 mg as an intravenous bolus. The Xylocaine is labeled 20 mg/mL. How many milliliters should be administered?

$$\frac{20 \text{ mg}}{1 \text{ mL}} = \frac{100 \text{ mg}}{x} = 5 \text{ mL}$$

12. Medication order: 50 mg/kg/day. Patient weight: 85.8 pounds. The patient will receive 1950 mg/day.  $85.8 = 39 \text{ kg} \times 50 = 1950 \text{ mg/day}$

13. Medication order: Amoxicillin 2.5 mL every 8 hours. Available is Amoxicillin 250 mg/5mL. The nurse will administer how many mg for the day?  $24/8 = \text{TID}$

$$\frac{250 \text{ mg}}{5 \text{ mL}} = \frac{x}{2.5 \text{ mL}} = 125 \text{ mg} \times 3 = 375 \text{ mg}$$

14. Medication order: Ondansetron 2 mg - 4 mg/kg/Q 4 hours po PRN nausea. The patient weighs 66 lbs. What is the minimum amount of medication in grams that can be administered every 4 hours?

$$66 / 2.2 = 30 \text{ kg} \times 2 \text{ mg} = 60 \text{ mg} \times \frac{1}{1000} = 60,000 \text{ g}$$

15. Medication order: 5 mL of normal saline is added to a vial of Lasix 20 mg/5 mL. How many milligrams of Lasix are in each millimeter of fluid?

$$\frac{20 \text{ mg}}{5 \text{ mL}} = \frac{x}{5 \text{ mL}} = \frac{100}{5} = 20 \text{ mg}$$