

Dosage Calculation Worksheet #4

1. Ordered is flucloxacillin 250mg IM. Available is 1 G in 10 mL. How much should the nurse administer in mL? 2.5mL

$$250/1000= 0.25 \times 10 = 2.5$$

2. Order: Administer 160 mg IV. Available is 100 mg/2 mL. How much should the nurse administer in mL? 3.2mL

$$160/100= 1.6 \times 2 = 3.2$$

3. Azulfidine 1.5 g has been ordered every 12 hrs. Available are 500mg tablets. How many tablets should the nurse administer per day? 6 tablets

$$1500/500=3 \times 2 = 6$$

4. Ergotrate maleate 200 mcg is ordered po daily. Available is 0.2 mg. How many tablets should the nurse administer? 1 tablet

$$200/1000=0.2$$

5. From 0700 to 1900 the nurse calculates the patient's total intravenous fluid intake as ? milliliters. An IV is infusing at 50 mL/hour. At 0900 the patient will receive IVPB of 125 mL for 30 minutes. What is the total amount in mL the patient will receive during this time? 700mL

$$50 \times 11.5 = 575 + 125 = 700\text{mL}$$

6. Solumedrol 1.5 mg/kg is ordered for a child weighing 42 lb. Solumedrol is available as 75 mg / 1 mL is available. How many mL must the nurse administer? 0.38mL

$$42/2.2= 19.1 \quad 1.5 \times 19.1= 28.65 \quad 28.65/75= 0.382 \rightarrow 0.38$$

7. Give patient 17.1 mg of dopamine in 223 mL of D5W to be infused at a rate of 17,221 mcg/hr. Calculate the flow rate in mL/hr. 224.6mL/hr

$$17,221/17,100 \times 223 = 224.57 \rightarrow 224.6$$

8. Calculate the IV flow rate for 0.2 L of D5W IV over 462 min. Infusion set has drop factor of 59 gtts/mL. What is the IV flow rate in gtts/min?

26gtts/min

$$200 \times 59 = 11,800 / 462 = 25.5 \rightarrow 26$$

9. Ordered Lasix 24 g IV push now. Available: 22,000,000 mcg in 12 mL. How much will the nurse draw up? 13.1mL

$$24,000,000 / 22,000,000 = 1.09 \times 12 = 13.08 \rightarrow 13.1$$

10. Calculate the IV flow rate for 392 mL of D5W IV over 582 min. Infusion set has drop factor of 74 gtts/mL. What is the IV flow rate in gtts/min?

50gtts/min

$$392 \times 74 = 29,008 / 582 = 49.8 \rightarrow 50$$

11. From 0700 to 1800 the nurse calculates the patient's total intravenous fluid intake as 1 milliliters. An IV is infusing at 100 mL/hour. At 0900 and 1500, the patient will receive IVPB of 75 mL for 30 minutes. What is the total amount the patient will receive during this time? 1050mL

$$9 \times 100 = 900 \quad 75 + 75 = 150 \quad 900 + 150 = 1050$$

12. Ordered 7 g of Amoxicillin. Amoxicillin is available as 0.016 kg per 20 mL. How much will the nurse draw up? 8.8mL

$$0.007/0.016 \times 20 = 8.75 \rightarrow 8.8$$

13. Potassium chloride is available as 0.016 kg per tablet. Potassium Chloride (K-Dur), 24,000,000 mcg, is ordered. How many tablets would the nurse administer? 1.5 tablets

$$24,000,000/16,000,000 = 1.5$$

14. Aggrastat at 23.8 mg in 129 mL is to be infused at 3 mcg/kg/hr in a patient who weighs 82 kg. At what flow rate in mL/hr will you set the pump? 1.3mL/hr

$$0.003 \times 82 = 0.246$$

$$0.246/23.8 \times 129 = 1.3$$

15. Administer 0.06 g of codeine po now. Available are 30 mg tablets. How many tablets should the nurse administer? 2 tablets

$$0.06/0.03 = 2$$