

9. From 0700 to 1600 the nurse calculates the patient's total intravenous fluid intake as $\boxed{1}$ milliliters. An IV is infusing at 150 mL/hour. At 1200, the patient will receive IVPB of 75 mL for 30 minutes. What is the total amount the patient will receive during this time?

1350

$$150 \times 8 = 2000$$

$$+ 75 \times 2$$



10. Administer 5 milligrams of acyclovir in 75 milliliters of normal saline over 15 minutes. The nurse will set the IV pump at $\boxed{1}$ mL/hour.

300

$$75 \times 4$$

$$60 \div 15 = 4$$

11. Phenytoin (Dilantin), ~~7,000,000~~ mcg PO, is ordered to be given through a nasogastric tube. Phenytoin is available as ~~5,000~~ mg / 18 mL. How much would the nurse administer? Round to a whole number.

25

$$\frac{7000 \times 18}{5}$$

12. Solumedrol 1.5 mg/kg is ordered for a patient weighing 74.8 lb. Solumedrol is available as 125 mg / 2mL. How many mL should the nurse administer?

0.82

$$1.5 \times (74.8 \div 2.2) = 51 \text{ mg}$$

$$\frac{51 \times 2}{125}$$

13. Give patient 24.4 mg of dopamine in 363 mL of D5W to be infused at a rate of 9,818 mcg/hr. Calculate the flow rate in mL/hr. ($363/24.4$)

146.1

9.818 mg

14. Give patient 10.1 mg of dopamine in 251 mL of D5W to be infused at a rate of 6 mg/hr. Calculate the flow rate in mL/hr.

149.1

$$251 / (10.1 / 6)$$

2500000

15. Ordered Lasix 12,000,000 mcg IV push now. Available: 0.025 kg in 15 mL. How much will the nurse draw up?

7.2

$$\frac{0.025 \text{ kg} \times 1000 \text{ mg/kg}}{12,000,000 \text{ mcg}} \times 15$$