

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	OXYTOCIC HORMONES	IT IS USE TO INDUCE LABOR AND CONTROL BLEEDING	0.5-40 UNITS	<u>Y</u> N	NaCl 0.9%	HYPOTENTION, FETAL HERAT RATE DECELERATION, NAUSEA, VOMITING, HEADACHE.	1. CONTINUE TO MONITOR MATERNAL BLOOD PRESSURE. 2. IF CONTRACTIONS BECOMES TOO FREQUENT, DECREASE INFUSION OR STOP MEDICATION. 3. START LOW AND GRADUALLY INCREASE INFUSION 4. ELECTRONICALLY MONITOR THE UTERINE ACTIVITY AND FETAL HEART RATE.
Magnesium Sulfate	ANTIDYSRHYTHMICS	USE TO TREAT PRECLAMPSIA IN PREGNANCIES	4G/ML 5G/ML	<u>Y</u> N	250ML NS/DSW	CIRCULATORY COLLAPSE, RESPIRATORY PARALYSIS, FLUSHING, DROWSINESS, INCREASED SWEATING	1. DO NOT USE ON PATIENT WHO HAS HIGH BLOOD PRESSURE 2. HYPERMAGNESEMIA MAY OCCUR 3. MONITOR RENAL FUNCTION 4. MAGNESIUM SULFATE IS SAFE FOR USE WHILE BREAST FEEDING
Meperidine	NARCOTICS	IT RELIEVES MODERATE TO SEVERE PAIN	25MG-100MG/ML	<u>Y</u> N		LIGHHEADEDNESS, HEADACHE, NAUSEA, CRAMPS	1. THIS MEDICATION CAN CAUSE IRREGULAR MENSTRUATION 2. REPORT ANY CHANGES IN HEARTBEAT 3. REPORT HAND TREMORS TO DOCTOR 4. REPORT ANY SLOW OR DIFFICULTY IN BREATHING
Promethazine	ANTI-HISTAMINE	IT IS USED TO	12.5	<u>Y</u>	25 ML NAACL 0.9%	FAST HEART	1. REPORT ANY BLACK TARRY STOOLS

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

	MINE	TREAT ALLERGY	MG-25MG	N		BEAT, HIGH FEVER, INCREASED SWEATING.	2.REPORT ANY BLEEDING GUMS, OR NOSE 3.THIA MEDICATION CAN CAUSE SLIGHT CONFUSION 4.IT CAN INCREASE HYPER ACTIVITY AND EXCITABILITY
Calcium Gluconate	ANTIDOTES	IT IS USED TO TREAT CALCIUM DEFICIENCIES	100MG/ML(10%)	<u>Y</u> N		DRY MOUTH, HEADCAHE, HOT FLASHES, DIARRHEA, CONSTIPATION	1.CALCIUM LELES NEED TO BE CHECKED 2.DO NOT STOP, START OR CHANGE DOSAGE BEFOR TALKING WITH HCP 3.IT CAN RESULT IN HYPERCALCEMIA 4.CARDIAC ARREST CAN OOCUR

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	VITAMIN	IT IS USED TO PREVENT BLEEDING	90MCG/PO	<u>Y</u> N		PAIN, SWELLING, TASTE CHANGES, SHORTNESS OF BREATH	1.IT CAN CAUSE UNUSUAL BLEEDING OR BRUIING 2.THIS MEDICATION CAN INTERACT WITH WARFARIN 3.REPORT ANY RASH, FEVER OR SWELLING OF FACE 4.IT CAN CAUSE TROUBLE BREATHING
Erythromycin Ophthalmic Ointment	MACROLIDE ANTIBIOTICS	IT TREATS BACTERIA INFECTION IN EYES		<u>Y</u> N		REDNESS, ITCHING, STINGING/BURNING OF EYES	1.VISION MAY BE BLURRY 2.DO NOT USE WITH OTHER OINTMENT 3.REPORT ANY RASH, FEVER AND HIVES 4.YOU MAY FEEL A STING OR BURNING IN EYES

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

Engerix B	HEPATITIS B VAACCINE	UES TO PREVENT HEPATITIS B	10MCG/ ML	Y N		CHEST PAIN, NAUSEA, PAIN IN ARMS, JAW, VOMITING	1.DO NOT GIVE IF ALLERGIC TO YEAST 2.IT CAN CAUSE SUDDEN NUMBNESS 3.WEAKNESS IN ARMS AND LEGS ARE COMMON 4.REPORT INABILITY TO MOVE ARMS AND LEGS
Hepatitis B Immune Globulin	ANTIVIRALS	USE TO PREVENT HEPATITIS FROM REOCCURING	0.06ML /KG	Y N		HIVES, SWEATING, FAINTNESS, LIGHTHEADEN ESS.	1.IT INCREASES RISK OF BLOOD CLOT 2.IT CAN CAUSE BLURRED VISION 3.IT MAY RESULT IN UNSUAL TIREDNESS OR WEAKNESS 4.CONFUSION MAY OCCUR
				Y N			1. 2. 3. 4.