

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
N/A	Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input type="checkbox"/>	N/A	N/A	N/A

Student Name: Naureen Doctolero		Unit: South 8	Patient Initials: CP		Date: 11/3/2020	Allergies: NKDA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Glimepiride	Sulfonylurea	Type 2 Diabetes mellitus	2 mg, PO	<input style="width: 100px; height: 20px;" type="text"/> here to enter text. Click here to enter text.		Dizziness, burred vision, fast heartbeat, hives, skin rash, puffiness or swelling of the eyelids, shortness of breath, wheezing	1. Take with breakfast or the first main meal of the day. Take with a full glass of water. 2. Avoid exposure to sunlight or tanning beds. Wear protective clothing and use sunscreen when you are outdoors. 3. Do not drive or perform hazardous tasks if glimepiride makes you drowsy, light-headed or dizzy. Avoid alcohol. 4. Make sure you eat regular meal and always carry a reliable source of glucose with you. Be alert for the symptoms of hypoglycemia (such as dizziness, headache, sweating, and shaking).
Metformin	Biguanide	Type 2 Diabetes	500 mg, PO	<input style="width: 100px; height: 20px;" type="text"/> here to enter text. Click here to enter text.		Diarrhea, nausea, flatulence, indigestion, headache, lack of energy, taste	1. Take with meal to reduce the risk of stomach upset. Once- daily dosages should be taken with the evening meal. 2. Swallow slow release tablets whole; do not crush, break, or chew.

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Naureen Doctolero		Unit: South 8	Patient Initials: CP	Date: 11/3/2020	Allergies: NKDA		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
						disturbance	3. Splitting dosages throughout the day (rather than taking a single dose) may improve gastric side effects such as nausea, diarrhea, and indigestion. 4. Do not drink large amount of alcohol or drink it daily while taking metformin because this may increase your risk of lactic acidosis.
Carvedilol	Beta blocker	High blood pressure, heart failure	25 mg, PO	<input type="text"/> here to enter text. Click here to enter text.		Hyperglycemia, angina pectoris, hypotension, nausea, orthostatic hypotension, rales, visual disturbance, vomiting	1. Take with food to reduce the risk of orthostatic hypotension. 2. Get up slowly from a sitting or lying down position to standing as carvedilol may cause a sudden drop in blood pressure. 3. Contact lens wearers may notice carvedilol decreases the natural lubrication of the eye; lubricant eye drops may be needed. 4. The dosage needs to be individualized and it may take some time to establish the correct dosage.
Lisinopril	Angiotensin converting	High blood pressure	20 mg, PO	<input type="text"/>	Click here to enter text.	Light headed feeling, fever,	1. Ensure adequate hydration before starting lisinopril.

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Naureen Doctolero		Unit: South 8	Patient Initials: CP	Date: 11/3/2020	Allergies: NKDA		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
	enzyme inhibitor (ACE inhibitor)	(hypertension)		Click here to enter text.		sore throat, cough, chest pain	2. Can be taken without regard to meals. 3. Lisinopril may make you feel dizzy, especially in the first few weeks of therapy and when going from a sitting or lying down position to standing. Always take your time when standing up. 4. Do not use salt substitutes or potassium supplements while taking lisinopril, unless recommended by your doctor.
Doxycycline	Tetracycline antibiotic	Bacterial infections	100 mg, PO	<input style="width: 100px; height: 20px;" type="text"/> here to enter text. Click here to enter text.		Difficult/painful swallowing, brown/gray tooth discoloration, unusual tiredness, abdominal pain, nausea, vomiting	1. Take doxycycline as directed. Some branded and generic versions of doxycycline need to be taken one hour prior to or two hours after meals. 2. Ensure you maintain hydration while taking doxycycline. This may help reduce the incidence of gastrointestinal side effects. 3. Avoid excessive sun exposure while receiving doxycycline. Seek medical advice if skin redness or skin eruptions develop. 4. A course of doxycycline may increase the risk of vaginal candidiasis in women. Talk to your doctor or pharmacist about

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Naureen Doctolero		Unit: South 8	Patient Initials: CP	Date: 11/3/2020	Allergies: NKDA		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							treatments if you develop symptoms of vaginal candidiasis.
Enoxaparin	Low molecular weight heparin	Deep vein thrombosis, pulmonary embolism	40 mg, SubQ	<input type="text"/> here to enter text. Click here to enter text.		Nausea, diarrhea, anemia, confusion, pain, bruising, lightheadedness, sudden weakness	<ol style="list-style-type: none"> 1. Alternate injection sites between left and right anterolateral and left and right posterolateral abdominal wall. 2. Do not expel the air bubble from the prefilled syringes before injection, to avoid the loss of drug. 3. To minimize bruising, do not rub the injection site after completion of the injection. 4. Instruct patient if taking a blood thinner, be sure to get instruction about how to take medicine safely. Blood thinners can cause serious bleeding problems.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	<input type="text"/> here to enter text. Click here to enter text.		Click here to enter text.	<ol style="list-style-type: none"> 1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	<input type="text"/> here to enter text. Click here to enter text.		Click here to enter text.	<ol style="list-style-type: none"> 1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text.

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Naureen Doctolero		Unit: South 8	Patient Initials: CP	Date: 11/3/2020	Allergies: NKDA		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text.

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Naureen Doctolero		Unit: South 8	Patient Initials: CP		Date: 11/3/2020	Allergies: NKDA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.