


Past Surgeries



Hysterectomy



74/ Female

Allergies: Penicillin, Lisinopril

Increase Confusion/ new onset of combativeness

Minor neurocognitive disorder
Alzheimer's vs vascular type

High WBC 11.8
 Low H&H 10/30
 Low magnesium 2.1
 High BNP 250
 Low pH 7.34
 High PCO2 50
 High HCO3 27
 Low Po2 88
 UA positive for leukocyte esterase, nitrites, ketones, crystals, cast, glucose, WBC, RBC, and bacteria.

Dehydration

Lactated Ringers

Isotonic/ hyperkalemia

Alzheimer's

Quetiapine

Anti-psychotics/ cough

Arthritis

ASA

Salicylates/ Rash

Enoxaparin

Anticoagulant/ site reaction

Protonix/ Pantoprazole

Antituler Agent/ abdominal pain

Geriatric Specialist

No Procedures or diagnostic studies at this time

Confusion, crying, fearful, agitated, combative, irregular labored breathing, decreased breath sounds, disoriented, weak

Nursing Interventions for Confusion:

1. I will make sure I explain everything I am doing to prevent any further confusion to the patient and promote a calm environment
2. Social Interaction is known to help with confusion so I will encourage the patient to interact with the hospital staff as well as her family

Discharge Planning: I will make sure the patient's caregiver knows the importance of a schedule adequate rest periods When caring for an Alzheimer's patient

Nursing Interventions weakness: Impaired physical mobility

1. I will look into the patients daily nutritional status and make sure they are getting adequate nutrition since that corallites to energy and an overall healthy body. I will make adjustments upon the results I find.
2. I will do ROM exercises with the patient 3x a day a little longer each time to help build her stamina

I will teach the patient some exercises she can do while at home to help build muscle. For example, use canned food as weights and do arm exercises while sitting down for safety.