

Instructional Module 2 Adult/Geriatric Medication Worksheet

Jesus Carrasco

1/13/2021

Student Name: _____

Date: _____

Allergies: Always Assess Before Administering Any Medication

Insulin Lispro "Humalog" Can be mixture: LA/SA	Short-Acting Insulin Pancreatic Faster-shorter life	Controls Hyperglycemia Anti-diabetic Hormones	Injection: Sub-Q Stomach, thighs, buttocks, back of forearm	Sliding Scale Insulin 1-10 hours of life Humalog Pen- Discard after 10 days 1 unit/0.01 mL 10mL syr. 100units/mL	Hypoglycemia Allergic Rxns: Rash, itching, swelling, difficulty breathing Heart Failure	1. Assess glucose levels prior to administration 2. Assess Lab: Potassium levels can cause cramps, irregular heartbeat 3. May cause dizziness, drowsiness, blurred vision 4. Have food tray available within 15 minutes
Furosemide	Diuretic Antihypertensive "Water Pill"	Reduces fluid in the body (Increases urination) Decreases HTN	PO IV IM	Diluent: NS, D5W and LR 4mL single dose vial 40mg (10mg/mL)	Black Box Warning: Irregular Potassium levels can cause dysrhythmias Dehydration	1. M/C: Drowsiness, headache, blurry vision, ringing in ears, hypersensitivity to sun 2. Can affect Blood glucose 3. Assess: Vitals (B/P) and Potassium Levels 4. Monitor: I/O, increase weight and edema
Vancomycin "Hydrochloride/ Vancocin"	Glycopeptide Antibiotic	Kills Bacteria in intestines	PO: Caps, solution IV infusion	Rate: 500mg/6hr IV 1g/200mL IV 250mg oral caps	Renal Failure Hearing loss	1. Assess: for LE edema, Vitals 2. Monitor: I/O, 3. Report: blood in urine/hearing loss 4. Assess Abdomen- May cause nausea, vomiting and pain.
Carafate	Antiulcer Dermatological	Protects and prevent ulcers	Nasogastric Oral	Oral: 1gm/10mL and 1gm tablets	Renal Failure in elderly or patients on dialysis Hyperglycemia	1. Monitor Blood glucose levels 2. Assess Abdomen- may cause constipation, bezoar formation, aluminum toxicity 3. Take on an empty stomach 4. Pt should not take antacids 30 pre or post drug.