

Student Name: Alyson Chauncey

Unit: OBS

Pt. Initials: N/A

Date: 2/22/2021

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: NKDA

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
N/A	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Pituitary Hormone, Uterine stimulant	Stimulates the uterus to produce contractions		Y N		N/V, blood pressure changes, cardiac dysrhythmias, water intoxication	1. FHR, UA, and fetal heart patterns need to be observed during the use of Oxytocin 2. Teach mother that this drug induces contractions during labor 3. Contraindicated in unfavorable fetal positions 4. Contraindicated with fetal distress if delivery is not imminent
Magnesium Sulfate	Anticonvulsant	Trace mineral that prevents seizures		Y N		Flushing, sweating, hypotension, hyporeflexia, CNS depression, respiratory tract paralysis	1. Serum magnesium needs to be drawn to ensure magnesium levels are within limits for dosage 2. Contraindicated in mothers with toxemia of pregnancy during the 2 hours before delivery 3. Renal impairment may lead to magnesium toxicity with this drug 4. Monitor for signs of toxicity with decreased respirations and cardiovascular hypotension
Meperidine	Analgesic, Opioid	Acts as pain reliever and sedative		Y N		Sweating, N/V, dizziness, sedation, hypotension, respiratory depression	1. Monitor for signs of respiratory depression 2. Black Box: has the potential for addiction/misuse 3. Prolonged use during pregnancy can result in neonatal opioid withdrawal syndrome 4. Contraindicated in acute or severe bronchial asthmatics without monitoring

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours**Allergies:** _____

Promethazine	Phenothiazine	Blocks histamine H1 receptors, sedative		Y N		Dermatitis, N/V, CNS depression, sedation, dizziness, prolonged QT interval	1. Contraindicated in patients with lower respiratory tract symptoms 2. Contraindicated in patients with asthma 3. Can cause prolonged QT interval so patients with cardiovascular disease need monitoring 4. Contraindicated in SQ administration
Calcium Gluconate	Calcium Supplement	Supplemental calcium		Y N		Abnormal taste in mouth, cardiac dysrhythmias, flatulence	1. Calcium is necessary for function of nervous, muscular, and skeletal systems 2. It is essential in nerve enzyme transmission 3. Contraindicated for IM or SQ routes 4. Contraindicated in elevated calcium levels

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	Vitamin K	Necessary for synthesis of prothrombin		Y N		Cardiac arrest, metabolic acidosis, CNS depression, Respiratory arrest	1. Use with caution in patients with preexisting liver disease 2. Hypersensitivity reactions may occur and can include shock 3. Monitor administration in premature infants due to benzyl alcohol in product can cause gasping syndrome 4. Max administration rate is 1mg/minute
Erythromycin Ophthalmic Ointment	Macrolide antibiotics	Stops the growth of bacteria		Y N		Eye irritation, burning of eye, redness	1. Contraindicated in patients hypersensitive or allergic to erythromycin 2. Used to prevent the growth of microorganisms 3. Avoid contaminating applicator tip 4. For ophthalmic use only
Engerix B	Vaccine	Vaccination for Hep B		Y N		Injection site reaction,	1. Non-infectious recombinant DNA hepatitis B 2. Contraindicated in neonates less than 2000 g at birth until 1 month of age or discharge

Student Name: _Alyson Chauncey_____

Unit: _OBS_____

Pt. Initials: _N/A_____

Date: _2/22/2021_____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _NKDA_____

						Nausea, fever	3. Apnea has been reported in premature infants 4. Administer vaccine within 12 hours of birth
Hepatitis B Immune Globulin	Immune Serum	Plasma product for immune response		Y N		Redness at injection site, N/ V, HA	1. Redness at site is most common side effect 2. Do not shake vial 3. Avoid live vaccines for at least 3 months after this drug is given 4. Human plasma always carries a risk for transmitting infectious agents