

Student Name: ASHLEE GUZZMAN

Date: 3/7/21

Allergies: Penicillins, Codeine, Meperidine

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
LACTATED RINGER 125/ml	<u>Iso</u> tonic/Hypotonic/Hypertonic	- IFS to replace fluid & electrolytes.	- POTASSIUM - acid base imbalance	- contraindicated to patients w/ hypersensitivity to corn products - severe liver disease

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Pantoprazole	- Proton pump inhibitors	- Antulcer agents	40mg PO Daily	<input checked="" type="checkbox"/> Y		Severe hepatic impairment, pruritus, flatulence, generalized edema	1. Advise patient that foods may cause increase in GI irritation. 2. Advise patient to report onset of black, tarry stools. 3. Instruct patient to notify any abdominal cramping to notify if renal products are being taken. 4. Instruct patient to notify if renal products are being taken.
Donepezil	Acetylcholinesterase improving cholinergic function	- Lessen some of dementia - enhances cognition	10mg PO Daily	<input checked="" type="checkbox"/> Y		depression, syncope, hypertension, frequent urination, fibrillation	1. Have patient with non-said focus since they will be dizzy. 2. Alert patient to get help from any nurse to get out of bed. 3. I will monitor my patient to always take her medication on time. 4. Inform patient we will see results within a week.
Duloxetine	Anti-depressant	major depressive disorder, fibromyalgia	30mg PO Daily	<input checked="" type="checkbox"/> Y		BLACKBOX warning of increase risk of suicidality, seizures, dysgeusia	1. Watch patient constantly & check for any unusual behavior. 2. Monitor patient for hematoma. 3. Monitor for hepatic failure so will have patient do bilirubin lab. 4. Put catheter on patient for uric acid retention.
Montelukast	Leukotriene receptor antagonist	decreases severity of allergic rhinitis	10mg PO Daily	<input checked="" type="checkbox"/> Y		Suicidal behavior thoughts, memory impairment, Steven Johnson syndrome, abdominal pain, liver enzymes	1. Have family to be alert for emergency of anxiety/panic attack. 2. Patient should carry rapid acting therapy for bronchospasm discontinuous. 3. Instruct patient to not discontinue med without consulting HCP. 4. Instruct patient to take medication daily in evening or 2hr prior exercise.
Sodium chloride	Normal Saline	prevent IV catheters from becoming blocked, remove any med left	10ml IV 12hrs	<input checked="" type="checkbox"/> Y		allergic reaction, urticaria at the injection site, redness, tightness at chest, swelling	1. Check compatibility of primary fluid with normal saline. 2. I will look to see any allergic reaction is occurring. 3. Check IV site regularly. 4. I will see that pump is pumping at the correct rate.