

Primary IV Fluid and Infusion Rate (mL/hr)	Circle IVF Type	Rational for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
NS 150 mL/hour	€ Isotonic € Hypotonic € Hypertonic	Maintenance Fluid	*CMP	*congestive heart failure *renal impairment *liver cirrhosis *sodium retention

Student Name: Bergan Gunter	Unit:	Patient Initials:	Date: 2/23/2021	Allergies:			
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP-List diluent solution, volume, and rate of administration IVPB-List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications , ETS.)
Enoxaparin (Lovenox)	Anticoagulant	Blood thinner	*SubQ: 30mg/0.3mL	yes		*Black box warning: epidural or spinal hematomas, which may result in long-term or permanent paralysis *Edema *Hematoma (bruise) *Confusion *Pain *Dyspnea (difficult or labored breathing)	1. Rotate injection sites 2. Patient should lie down during injection. 3. Tell patient to report signs/symptoms of bleeding. 4. Tell patient to avoid taking other anticoagulants such as aspirin while taking enoxaparin. 5. assess platelet count. 6. Tell patient not to rub the

						*Fever	injection site to avoid bruising.
Lasix (Furosemide)	Loop Diuretic	*Treatment of hypercalcemia	*IVP: 20mg/2mL	yes	Slowly over 1-2 min	<p>*Hyperuricemia (too much uric acid in your blood)</p> <p>*Hypomagnesemia (serum magnesium concentration <1.8 mg/dL)</p> <p>*Loss of appetite</p> <p>*Spasm of bladder</p> <p>*Hypotension</p> <p>*Pancreatitis</p>	<p>1. Drug causes sun sensitivity</p> <p>2. Instruct patient to report unusual bleeding/bruising or signs/symptoms of hypotension, infection, pancreatitis, or ototoxicity (tinnitus, hearing impairment)</p> <p>3. Instruct patient to eat high-potassium foods during drug therapy, as directed by healthcare professional.</p> <p>4. Patient should not drink alcohol while taking this drug.</p> <p>5. Monitor BP, serum electrolytes, calcium, potassium, BUN and more</p>
Protonix (Pantoprazole)	*Gastric Acid Secretion Inhibitor *Proton Pump	Decreases the amount of acid in the stomach which	*IVP: 40mg / 10mL	yes	Over 1-2 min *40 mg reconstituted with 10mL NS	<p>*Abdominal pain</p> <p>*Flatulence</p> <p>*C diff</p> <p>*Tubulointerstiti</p>	<p>1. Teach patient to report diarrhea that does not improve</p> <p>2. Side effects may include headache, diarrhea,</p>

	Inhibitor	relieves symptoms of heartburn, difficulty swallowing , and persistent cough				al nephritis (inflammation that affects the tubules of the kidneys and the tissues that surround them; interstitial tissue)	nausea, abdominal pain, vomiting, flatulence, dizziness, arthralgia, rash, and fever 3. Tell patient to report symptoms of Tubulointerstitial Nephritis (excessive urination, fever and or rash) 4. Have patient report symptoms of tremors, weakness, and exhaustion (hypomagnesemia symptoms)