

Student Name: \_\_\_\_\_

Unit: \_\_\_\_\_

Pt. Initials: \_\_\_\_\_

Date: \_\_\_\_\_

**Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Uterine stimulant	Promotes contractions.		Y N	Dilute 10 units in 1000 mL of NS or LR.	N/V, fetal bradycardia, increased BP	1. Give bolus of 500ml if tachysystole 2. Stop oxytocin with tachysystole or abnormal FHR pattern 3. Stop with category 2 or 3. 4.
Magnesium Sulfate	Anticonvulsant	Prevent seizures		Y N	Dilute with D5W or NS to a concentration of 20% or less.	Flushing, sweating, hypotension, CNS depression	1. Monitor BP closely. 2. Place resuscitation equipment in the room. 3. Calcium gluconate is the antidote. 4. Monitor for RR of at least 12.
Meperidine	Opioid	For pain management		Y N	Can be given PO, SQ, and IM.	Sweating, N/V, Dizziness, Lightheaded, constipation.	1. Potential for addiction, and abuse. 2. Report s/s of resp depression. 3. Naloxone is the antidote. 4. Teach about constipation.
Promethazine	Antihistamine	For N/V		Y N	Can be given SQ, PO, IM, or IV.	Dizziness, sedated, low convulsive threshold.	1. Photosensitivity, avoid sunlight, use sunscreen. 2. Avoid activities requiring mental alertness until effects are known. 3. No MAO inhibitors or CNS depressants while on promethazine. 4. Caution with patients with hx of seizures.

Student Name: \_\_\_\_\_

Unit: \_\_\_\_\_

Pt. Initials: \_\_\_\_\_

Date: \_\_\_\_\_

**Newborn Medication Worksheet – Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

Calcium Gluconate	Calcium supplement	Calcium replacement		Y N		Strange taste in mouth, constipation, flatulence, swollen abdomen	<ol style="list-style-type: none"> <li>Warn pt for chalk taste, tingling, and sensation of heat waves.</li> <li>Patient should take oral form with food or liquid.</li> <li>Chew oral tablets and follow with water.</li> <li>Monitor for s/s of bradycardia abd hypotension</li> </ol>
-------------------	--------------------	---------------------	--	--------	--	---	---

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	Vitamin K	Prevents Bleeding		Y N		CNS depression, cyanosis, dyspnea, flushing.	<ol style="list-style-type: none"> <li>Administer 1mg/minute when IV.</li> <li>Teach pt to report rashes.</li> <li>Avoid increases in Vit K food such as green veggies, pork, and beef liver.</li> <li>Black box warning: fatalities have occurred immediately after IV therapy.</li> </ol>
Erythromycin Ophthalmic Ointment	Antibiotic	Prevents eye infection		Y N		Stinging, burning, blurred vision	<ol style="list-style-type: none"> <li>Blurred vision after application is normal.</li> <li>Applied to infants prophylactically</li> <li></li> <li></li> </ol>
Engerix B	Vaccine	Prevents Hep B Infection		Y N		Low fever, sore arm.	<ol style="list-style-type: none"> <li>First dose within 24 hours of birth.</li> <li>Second dose is at one month, and third dose is at 6 months.</li> <li></li> <li></li> </ol>
Hepatitis B Immune	Immune serum	Prevents Hep B Infection		Y N		Erythema at injection site, HA, lowered	<ol style="list-style-type: none"> <li>Works by immediately providing protection from Hep B infection, not long term.</li> <li>Avoid live vaccines for 3 months after</li> </ol>

Student Name: \_\_\_\_\_

Unit: \_\_\_\_\_

Pt. Initials: \_\_\_\_\_

Date: \_\_\_\_\_

**Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours**

**Allergies:** \_\_\_\_\_

Globulin						WBC.	receiving this, except for hep B vaccine. 3. Watch patient for 15 minutes after administering incase of reaction. 4.
				Y N			1. 2. 3. 4.