

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

| Primary IV Fluid and Infusion Rate (ml/hr) | Circle IVF Type | Rationale for IVF | Lab Values to Assess Related to IVF | Contraindications/Complications |
|--|---------------------------------|-------------------|-------------------------------------|---------------------------------|
| | Isotonic/ Hypotonic/ Hypertonic | | | |

| Generic Name | Pharmacologic Classification | Therapeutic Reason | Dose, Route & Schedule | Correct Dose? If not, what is correct dose? | IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give | Adverse Effects | Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.) |
|-------------------|------------------------------|--------------------|---|---|---|---|--|
| Oxytocin | Exogenous hormones | Oxytocics | 10 U/1 mL | Y N | n/a | Postpartum hemorrhage, uterine rupture, abruptio placentae, hematoma, hypotension, tachycardia, N/V | 1. Monitor and record uterine contractions, HR, BP, intrauterine pressure, fetal HR, and character of blood loss at least every 15 minutes. 2. Discontinue immediately if uterine hyperactivity or fetal distress occurs. 3. Use cautiously in pregnant or breastfeeding women. 4. All patients receiving oxytocin IV must be under continuous observation. |
| Magnesium Sulfate | Magnesium salts | Laxative | INJ - 500 mg/ 1 mL IV - 40 mg/1 mL, 80 mg/1 mL Powder for solution - 100% | Y N | | Flushing, sweating, hypotension, shock, hyporeflexia | 1. This drug may cause hypotension, hyporeflexia or CNS depression. 2. Use cautiously in patients with sodium restrictions because drug contains sodium. 3. Monitor electrolyte levels during use. 4. Overdose s/s include blurred or double vision, coma, dizziness, syncope, drowsiness, bradycardia, and dyspnea. |
| Meperidine | Analgesic/ opioid | Analgesic | INJ - 25, 50, 75, or | Y N | | Myoclonus, seizure, shock, orthostatic | 1. Be aware this drug can cause significant respiratory depression. 2. Contraindicated with MAOI therapy or |

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

| | | | | | | | |
|-------------------|---------------|---------------|---|------------|-----|---|---|
| | | | 100 mg/ mL IV - 10 mg/1 mL PO - 50, 100 mg | | | hypotension, respiratory depression, N/V, sweating | recent use within 14 days. 3.Report severe constipation. 4.Avoid activities requiring mental alertness. |
| Promethazine | Phenothiazine | Antihistamine | INJ - 25, 50 mg/1 mL PO - 12.5, 25, 50 mg Rectal supposi- tory - 12.5, 25, 50 mg | Y N | n/a | Urticaria, N/V, phototoxicity, lowered convulsive threshold, CNS depression, neuroleptic malignant syndrome | 1.Advise patient to avoid prolonged exposure to sunlight. 2.Avoid drinking alcohol while taking this drug. 3.Do not take with MAO inhibitors or CNS depressants with this drug. 4.Report sweating, fever, stupor, unstable BP, muscular rigidity, and autonomic dysfunction. |
| Calcium Gluconate | Calcium salts | Antidote | IV - 100 mg/1 mL PO - 500 mg | Y N | | Constipation, abnormal taste in mouth, flatulence, hypotension | 1.May cause tingling sensations, chalky taste, a sense of oppression, or heat waves. 2.Patient should take oral powder with food or liquid. 3.Monitor serum calcium levels. 4.If in tablet form, swallow or chew thoroughly, followed by a glass of water. |

| | | | | |
|--|-----------------|-------------------|-------------------------------------|---------------------------------|
| Primary IV Fluid and Infusion Rate (ml/hr) | Circle IVF Type | Rationale for IVF | Lab Values to Assess Related to IVF | Contraindications/Complications |
|--|-----------------|-------------------|-------------------------------------|---------------------------------|

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

| | | Isotonic/ Hypotonic/ Hypertonic | | | | | |
|----------------------------------|----------------------------------|---------------------------------|--|---|---|--|---|
| Generic Name | Pharmacologic Classification | Therapeutic Reason | Dose, Route & Schedule | Correct Dose? If not, what is correct dose? | IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give | Adverse Effects | Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.) |
| Phytonadione | Vitamins | Vitamin K | PO - 2.5 to 10 mg IV - 2.5 to 10 mg | Y N | | Cardiac arrest, shock, metabolic acidosis, respiratory arrest | 1. Tell patient to report rashes. 2. Report feeling weak, tingly, lightheaded, chest pain, swelling in your face, or cold sweats. 3. Reversal of vitamin K is warfarin. 4. Protect this from the light. |
| Erythromycin Ophthalmic Ointment | Macrolide antibiotics | Antibiotic | Neonates - 10 mg/kg/dose PO q 6 | Y N | n/a | Diarrhea, loss of appetite, stomach cramps, seizure, ototoxicity, vomiting | 1. Avoid contact with eyes and all mucus membranes. 2. Use a tissue to wipe excess from eyelashes. 3. Blurred vision can occur for a short time. 4. Wash hands before applying. |
| Engerix B | Vaccine | Prophylaxis | INJ - 20 mcg/mL | Y N | n/a | Pain, soreness, redness, swelling, fever, headache, dizziness, injection site reaction | 1. Administer the first dose to all newborns before hospital discharge. 2. Unvaccinated children should complete a 3 dose series. 3. Administer in the deltoid for adults and older children and in younger children give in the thigh. 4. Never given IV/intradermal. |
| Hepatitis B Immune Globulin | Disease specific immunoglobulins | Antiviral | IM slowly | Y N | n/a | Allergic reaction, itching or hives, swelling of the face, chest pain | 1. This is refrigerated (36 to 46 degrees F). 2. Immune globulins may diminish the therapeutic effect of vaccines. 3. Use with caution in patients with |

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

| | | | | | | | |
|--|--|--|--|--------|--|--|---|
| | | | | | | or tightness, breathing problems, pain and tenderness at the injection site | thrombocytopenia or coagulation disorders. 4. IM injections may be contraindicated in patients with bleeding disorders. |
| | | | | Y N | | | 1. 2. 3. 4. |