

Maternal Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: _____

| Primary IV Fluid and Infusion Rate (ml/hr) | Circle IVF Type | Rationale for IVF | Lab Values to Assess Related to IVF | Contraindications/Complications |
|--|---------------------------------|-------------------|-------------------------------------|---------------------------------|
| | Isotonic/ Hypotonic/ Hypertonic | | | |

| Generic Name | Pharmacologic Classification | Therapeutic Reason | Dose, Route & Schedule | Correct Dose? If not, what is correct dose? | IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give | Adverse Effects | Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.) |
|-------------------|------------------------------|--|-----------------------------|---|---|--|--|
| Oxytocin | Oxytocic hormones | Induces labor or strengthen uterine contractions or to control bleeding after childbirth | Can be give IM or IV | Y N | | <ul style="list-style-type: none"> Slow/fast heart rate N/V Pounding in neck or ears Severe HA, blurred vision | <ol style="list-style-type: none"> Monitor fetal heart rate before, during, and after administration Monitor intrauterine pressure Assess cardiovascular system before, during, and after administration Review patients medical record prior to administration to be sure there are no contradictions |
| Magnesium Sulfate | Antidysrhythmics | Controls low blood levels of Mg, and prevent seizures in severe pre-eclampsia, eclampsia or toxemia of pregnancy | Can be given PO, IM, and IV | Y N | | <ul style="list-style-type: none"> Upset stomach Diarrhea Hypotension Increased sweating Contractions Breathing difficulties | <ol style="list-style-type: none"> Monitor fetal heart rate before, during, and after administration Assess respiratory system Assess cardiovascular system Inform patient to report any GI upset |
| Meperidine | Opiate (narcotic) analgesics | Treats moderate to severe pain | Can be given PO and IV | Y N | | <ul style="list-style-type: none"> Dizziness Drowsiness N/V Sweating | <ol style="list-style-type: none"> Monitor for serotonin syndrome Monitor fetal heart rate before, during, and after administration Monitor for respiratory depression Assess pain level before, during, after admin |
| Promethazine | Antiemetic | Treatment of nausea, vomiting | Can be IV or PO | Y N | Give over 3-5 mins | BBW – can cause fatal respiratory depression in pts under 2 years old. – CNS depression, dizziness, | <ol style="list-style-type: none"> May cause photosensitivity Report s/s of neuroleptic malignant syndrome (sweating, fever, stupor, unstable BP, muscular rigidity) |

Newborn Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: _____

| | | | | | | | |
|-------------------|--------------------|--|-----------------|--------|--|---|--|
| | | | | | | sedation, somnolence, prolonged QT interval | 3. Do not take MAOI's with this medication 4. Report s/s of respiratory depression |
| Calcium Gluconate | Calcium Supplement | Treatment of hypocalcemia due to rapid growth or pregnancy | Can be IV or PO | Y N | | <ul style="list-style-type: none"> Dysuria Swelling Weight gain Lightheadedness Bradycardia or dysrhythmias N/V Constipation | 1. Report s/s of bradycardia or dysrhythmias 2. My cause hypotension and syncope so change positions and get up slowly and with help 3. Monitor serum calcium levels 4. If taking PO, take with food or full glass of water |

| Primary IV Fluid and Infusion Rate (ml/hr) | Circle IVF Type | Rationale for IVF | Lab Values to Assess Related to IVF | Contraindications/Complications |
|--|---------------------------------|-------------------|-------------------------------------|---------------------------------|
| | Isotonic/ Hypotonic/ Hypertonic | | | |

| Generic Name | Pharmacologic Classification | Therapeutic Reason | Dose, Route & Schedule | Correct Dose? If not, what is correct dose? | IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give | Adverse Effects | Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.) |
|----------------------------------|------------------------------|-------------------------------|----------------------------|---|---|--|---|
| Phytonadione | Nutriceutical | Vitamin K | Can be IV, PO, SubQ, or IM | Y N | Dilute in NS or D5W in minimum of 50 mL over 20 mins Do not exceed 1mg/min | BBW – Risk of fatal reaction like anaphylaxis <ul style="list-style-type: none"> Cyanosis Dizziness Dysgeusia Dyspnea Flushing | 1. Monitor for signs of severe reaction or anaphylaxis 2. Report rash 3. Monitor prothrombin time 4. Monitor for respiratory arrest or breathing difficulties |
| Erythromycin Ophthalmic Ointment | Antibiotic | Treat bacterial eye infection | | Y N | | Urticaria, abdominal pain, diarrhea, loss of appetite, nausea, vomiting | 1. This drug may cause abdominal pain, loss of appetite, nausea, vomiting, or flu like symptoms. 2. Report signs of hepatic dysfunction. 3. Instruct patient to report severe diarrhea and consult PCP before taking anti-diarrhea medications. |

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

| | | | | | | | |
|-----------------------------|--------------|--|--|--------|--|---|---|
| | | | | | | | 4. May take on an empty stomach or right before meals. |
| Engerix B | Vaccine | Hep B vaccine | | Y N | | Injection site reaction, diarrhea, nausea, headache, fever, malaise | <ol style="list-style-type: none"> 1. This drug may cause diarrhea, nausea, asthenia, headache, fatigue, fever, or malaise. 2. Review dosing schedule with patient. 3. Call your doctor if you are feeling light-headed, like you might pass out. 4. Report a fever, and swollen glands. |
| Hepatitis B Immune Globulin | Immune serum | Perinatal exposure of infants born to HBsAg-positive mothers | | Y N | | Redness at injection site, nausea, vomiting, decreased WBC, ecchymosis, joint stiffness, headache | <ol style="list-style-type: none"> 1. Avoid live vaccines other than hepatitis B vaccine for 3 months after receiving this drug. 2. This drug may cause nausea, vomiting, itching, joint stiffness, malaise. 3. May interfere with some types of blood glucose monitoring systems and can result in falsely elevated glucose readings. 4. Call your doctor if you have fever, mouth sores, red or swollen gums. |
| | | | | Y N | | | <ol style="list-style-type: none"> 1. 2. 3. 4. |